antibodies

Datasheet for ABIN3088929 AGXT2 Protein (AA 42-514) (His tag)





Overview

Quantity:	1 mg
Target:	AGXT2
Protein Characteristics:	AA 42-514
Origin:	Human
Source:	Insect Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This AGXT2 protein is labelled with His tag.
Application:	Western Blotting (WB), SDS-PAGE (SDS), ELISA, Crystallization (Crys)

Product Details

Sequence:	LHTKPRMPPC DFMPERYQSL GYNRVLEIHK EHLSPVVTAY FQKPLLLHQG HMEWLFDAEG
	SRYLDFFSGI VTVSVGHCHP KVNAVAQKQL GRLWHTSTVF FHPPMHEYAE KLAALLPEPL
	KVIFLVNSGS EANELAMLMA RAHSNNIDII SFRGAYHGCS PYTLGLTNVG TYKMELPGGT
	GCQPTMCPDV FRGPWGGSHC RDSPVQTIRK CSCAPDCCQA KDQYIEQFKD TLSTSVAKSI
	AGFFAEPIQG VNGVVQYPKG FLKEAFELVR ARGGVCIADE VQTGFGRLGS HFWGFQTHDV
	LPDIVTMAKG IGNGFPMAAV ITTPEIAKSL AKCLQHFNTF GGNPMACAIG SAVLEVIKEE
	NLQENSQEVG TYMLLKFAKL RDEFEIVGDV RGKGLMIGIE MVQDKISCRP LPREEVNQIH
	EDCKHMGLLV GRGSIFSQTF RIAPSMCITK PEVDFAVEVF RSALTQHMER RAK
	Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a
	special request, please contact us.
Characteristics:	 Made in Germany - from design to production - by highly experienced protein experts. Human AGXT2 Protein (raised in Insect Cells) purified by multi-step, protein-specific process

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/4 | Product datasheet for ABIN3088929 | 09/11/2023 | Copyright antibodies-online. All rights reserved.

Product D	etails
-----------	--------

to ensure crystallization grade.

• State-of-the-art algorithm used for plasmid design (Gene synthesis).

experts in the lab will ensure that you receive a correctly folded protein.The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).When you order this made to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered. The concentration of our recombinant proteins is measured using the absorbance at 280nm. The protein saborbance will be measured in several dilutions and is measured against its apecific reference buffer. The concentration of proteins expressed in baculovirus infected SF9 insect cells: 1. In a first purification atep the protein is pacific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.Purification:Two step purification of proteins expressed in baculovirus infected SF9 insect cells: 1. In a first purification step, the protein is purification are subjected to second purification step through size exclusion chromatography. Elust fractions are analyzed by SDS-PACE and Western blot.Purification:>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.Sterility:0.22 µm filteredEndotoxin Level:Protein is endotxin free.<		This protein is a made to order protein and will be made for the first time for your order. Our
The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified. In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for oustom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization). When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experimened protein experts will do everything to make sure that you receive the protein you ordered. The concentration of our recombinant proteins is measured using the absorbance at 280nm. The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer. The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein. Unification are subjected SP9 insect cells: 1. In a first purification after, the protein is purified from the cleared cell ysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGEPurify:>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.Sterility:0.22 µm filteredEndotoxin Level:Protein is endotoxin free.Orade:Cystallography gradeTarget DetailisAOXT2		
made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified. In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization). When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experise will do everything to make sure that you receive the protein you ordered. The concentration of our recombinant proteins is measured using the absorbance at 280nm. The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer. The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.Purification:Two step purification step, the protein is purified from the cleared cell ysate using three different His-tag capture materials, high yield, EDTA resistant, or DTT resistant. Eluste fractions are analyzed by SDS-PAGE. 2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE 2. Protein containing fractions of the best purification are analyzed by SDS-PAGE 2. Protein containing fractions of the best purification are analyzed by SDS-PAGE 2. Protein containing fractions of the best purification are analyzed by SDS-PAGE 2. Protein containing fractions of the best purification are analyzed by SDS-PAGE and Western blot.Purity:>95 % as det		
cannot be expressed or purified. In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for oustom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization). When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered. The concentration of our recombinent proteins is measured using the absorbance at 280nm. The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer. The concentration of proteins expressed in baculovirus infected SF9 insect cells: 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE. 2. Protein containing fractions of the best purification are analyzed by SDS-PAGE and Western blot.Purity:>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.Sterility:0.22 µm filteredEndotxin Level:Protein is endotxin free.Grade:Crystallography gradeTarget DetailisAGXT2		
In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization). When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered. The concentration of our recombinant proteins is measured using the absorbance at 280nm. The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer. The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials. high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE. 2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.Purity:>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.Endotoxin Level:Protein is endotoxin free.Grade:Crystallography grade.Target DetailsCrystallography grade.		
(other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization). When you order this made to order protein you will only pay upon receival of the correctly foided protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered. The concentration of our recombinant proteins is measured using the absorbance at 280nm. The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer. The concentration of proteins expressed in baculovirus infected SF9 insect cells: 1. In a first purification of protein se pressed in baculovirus infected SF9 insect cells: 1. In a first purification step, the protein is purified from the cleared cell lysate using three different Histag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.Purity:>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.Sterility:0.22 µm filteredEndotoxin Level:Protein is endotoxin free.Grade:Crystallography gradeTarget DetailsTarget		
custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered. The concentration of our recombinant proteins is measured using the absorbance at 280nm. The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer. The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.Purification:Two step purification step, the protein is purified from the cleared cell lysate using three different His-tag capfure materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE. 2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE 2. Protein is endotoxin free.Purity:>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.Sterility:0.22 µm filteredEndotoxin Level:Crystallography gradeTarget DetailsCrystallography grade		
experiments or purification optimization). When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered. The concentration of our recombinant proteins is measured using the absorbance at 280nm. The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer. The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.Purification:Two step purification of proteins expressed in baculovirus infected SF9 insect cells: 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE. 2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.Purity:>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.Sterility:0.22 µm filteredEndotoxin Level:Protein is endotoxin free.Grade:Crystallography gradeTarget DetailsAGXT2		
folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered. The concentration of our recombinant proteins is measured using the absorbance at 280nm. The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer. The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.Purification:Two step purification of proteins expressed in baculovirus infected SF9 insect cells: 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE. 2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.Purity:>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.Sterility:0.22 µm filteredEndotoxin Level:Protein is endotoxin free.Grade:Crystallography gradeTarget DetailsAGXT2		experiments or purification optimization).
protein experts will do everything to make sure that you receive the protein you ordered. The concentration of our recombinant proteins is measured using the absorbance at 280nm. The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer. The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.Purification:Two step purification of proteins expressed in baculovirus infected SF9 insect cells: 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE. 2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE.Purity:>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.Sterility:0.22 µm filteredEndotoxin Level:Protein is endotoxin free.Grade:Crystallography gradeTarget DetailsAGXT2		When you order this made-to-order protein you will only pay upon receival of the correctly
The concentration of our recombinant proteins is measured using the absorbance at 280nm. The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer. The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.Purification:Two step purification of proteins expressed in baculovirus infected SF9 insect cells: 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE. 2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.Purity:>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.Sterility:0.22 µm filteredEndotoxin Level:Protein is endotoxin free.Grade:Crystallography gradeTarget DetailsA6XT2		folded protein. With no financial risk on your end you can rest assured that our experienced
The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer. The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.Purification:Two step purification of proteins expressed in baculovirus infected SF9 insect cells: 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE. 2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.Purity:>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.Sterility:0.22 µm filteredEndotoxin Level:Protein is endotoxin free.Grade:Crystallography gradeTarget DetailsAGXT2		protein experts will do everything to make sure that you receive the protein you ordered.
specific reference buffer. The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.Purification:Two step purification of proteins expressed in baculovirus infected SF9 insect cells: 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE. 2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.Purity:>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.Sterility:0.22 µm filteredEndotoxin Level:Protein is endotoxin free.Grade:Crystallography gradeTarget DetailsAGXT2		The concentration of our recombinant proteins is measured using the absorbance at 280nm.
The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.Purification:Two step purification of proteins expressed in baculovirus infected SF9 insect cells: 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.Purity:-Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.Purity:-95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.Sterility:0.22 µm filteredEndotoxin Level:Protein is endotoxin free.Grade:Crystallography gradeTarget DetailsAGXT2		The protein's absorbance will be measured in several dilutions and is measured against its
the Expasy's protparam tool to determine the absorption coefficient of each protein.Purification:Two step purification of proteins expressed in baculovirus infected SF9 insect cells:1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.Purity:>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.Sterility:0.22 µm filteredEndotoxin Level:Protein is endotoxin free.Grade:Crystallography gradeTarget DetailsAGXT2		specific reference buffer.
Purification:Two step purification of proteins expressed in baculovirus infected SF9 insect cells:1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGEPurity:>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.Sterility:0.22 µm filteredEndotoxin Level:Protein is endotoxin free.Grade:Crystallography gradeTarget DetailsAGXT2		The concentration of the protein is calculated using its specific absorption coefficient. We use
 In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot. Purity: >95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot. Sterility: 0.22 µm filtered Protein is endotoxin free. Grade: Crystallography grade Target Details Target: AGXT2 		the Expasy's protparam tool to determine the absorption coefficient of each protein.
different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.Purity:>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.Sterility:0.22 µm filteredEndotoxin Level:Protein is endotoxin free.Grade:Crystallography gradeTarget DetailsAGXT2	Purification:	Two step purification of proteins expressed in baculovirus infected SF9 insect cells:
fractions are analyzed by SDS-PAGE.2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.Purity:>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.Sterility:0.22 μm filteredEndotoxin Level:Protein is endotoxin free.Grade:Crystallography gradeTarget DetailsAGXT2		1. In a first purification step, the protein is purified from the cleared cell lysate using three
2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.Purity:>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.Sterility:0.22 µm filteredEndotoxin Level:Protein is endotoxin free.Grade:Crystallography gradeTarget DetailsAGXT2		
through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.Purity:>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.Sterility:0.22 µm filteredEndotoxin Level:Protein is endotoxin free.Grade:Crystallography gradeTarget DetailsAGXT2		
Purity:>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.Sterility:0.22 µm filteredEndotoxin Level:Protein is endotoxin free.Grade:Crystallography gradeTarget DetailsAGXT2		
Sterility:0.22 µm filteredEndotoxin Level:Protein is endotoxin free.Grade:Crystallography gradeTarget DetailsAGXT2		Western blot.
Endotoxin Level:Protein is endotoxin free.Grade:Crystallography gradeTarget DetailsAGXT2	Purity:	>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.
Grade: Crystallography grade Target Details AGXT2	Sterility:	0.22 µm filtered
Target Details Target: AGXT2	Endotoxin Level:	Protein is endotoxin free.
Target: AGXT2	Grade:	Crystallography grade
Target: AGXT2	Target Details	
		ΔGXT2
Alternative Name: AGXT2 (AGXT2 Products)		
	Alternative Name:	AGXT2 (AGXT2 Products)

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/4 | Product datasheet for ABIN3088929 | 09/11/2023 | Copyright antibodies-online. All rights reserved.

Target Details	
Background:	Can metabolize asymmetric dimethylarginine (ADMA) via transamination to alpha-keto-delta- (NN-dimethylguanidino) valeric acid (DMGV). ADMA is a potent inhibitor of nitric-oxide (NO) synthase, and this activity provides mechanism through which the kidney regulates blood pressure. {ECO:0000269 PubMed:20018850, ECO:0000269 PubMed:23023372, ECO:0000269 PubMed:24586340}.
Molecular Weight:	53.5 kDa Including tag.
UniProt:	Q9BYV1
Pathways:	Monocarboxylic Acid Catabolic Process
Application Details	
Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.
Comment:	In cases in which it is highly likely that the recombinant protein with the default tag will be insoluble our protein lab may suggest a higher molecular weight tag (e.g. GST-tag) instead to increase solubility. We will discuss all possible options with you in detail to assure that you receive your protein of interest.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	Unlimited (if stored properly)



Image 1. "Crystallography Grade" protein due to multi-step, protein-specific purification process

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 4/4 | Product datasheet for ABIN3088929 | 09/11/2023 | Copyright antibodies-online. All rights reserved.