

Datasheet for ABIN3089295
ARIH1 Protein (AA 1-557) (Strep Tag)



[Go to Product page](#)

1 Image

Overview

Quantity:	1 mg
Target:	ARIH1
Protein Characteristics:	AA 1-557
Origin:	Human
Source:	Tobacco (Nicotiana tabacum)
Protein Type:	Recombinant
Purification tag / Conjugate:	This ARIH1 protein is labelled with Strep Tag.
Application:	ELISA, SDS-PAGE (SDS), Western Blotting (WB)

Product Details

Sequence: MDSDEGYNYE FDEDEECSEE DSGAEEEEDE DDDEPDDDTL DLGEVELVEP GLGVGGERDG
LLCGETGGGG GSALGPGGGG GGGGGGGGGG PGHEQEEDYR YEVLTAEQIL QHMVECIREV
NEVIQNPATI TRILLSHFNW DKEKLMERYF DGNLEKLFAE CHVINPSKKS RTRQMNRSS
AQDMPQCICY LNYPN SYFTG LECGHKFCMQ CWSEYLTTKI MEEGMGQTIS CPAHGCDILV
DDNTVMRLIT DSKVKLKYQH LITNSFVECN RLLKWCPAPD CHHVVKVQYP DAKPVRCKCG
RQFCFNCGEN WHDPVKCKWL KKWIKKCCDD SETSNWIAAN TKECPKCHVT IEKDGGCNHM
VCRNQCKAE FCWVCLGPWE PHGSAWYNCN RYNEDDAKAA RDAQERSRAA LQRYLYFCNR
YMNHMQLRF EHKLYAQVKQ KMEEMQQHNM SWIEVQLFKK AVDVLQCQRA TLMYTYVFAF
YLKKNQSI FENNQADLEN ATEVLSGYLE RDISQDSLQD IKQKVQDKYR YCESRRRVLL
QHVHEGYEKD LWEYIED

Sequence without tag. The proposed Strep-Tag is based on experience s with the expression system, a different complexity of the protein could make another tag necessary. In case you

have a special request, please contact us.

Characteristics:

Key Benefits:

- Made in Germany - from design to production - by highly experienced protein experts.
- Protein expressed with ALiCE® and purified by multi-step, protein-specific process to ensure correct folding and modification.
- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from *Nicotiana tabacum* c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.
- During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.
- We use the Expasy's ProtParam tool to determine the absorption coefficient of each protein.

Purification:

Two step purification of proteins expressed in Almost Living Cell-Free Expression System (ALiCE®):

1. In a first purification step, the protein is purified from the cleared cell lysate using StrepTag capture material. Eluate fractions are analyzed by SDS-PAGE.

Product Details

2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

Purity: >80 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.

Endotoxin Level: Low Endotoxin less than 1 EU/mg (< 0.1 ng/mg)

Grade: Crystallography grade

Target Details

Target: ARIH1

Alternative Name: ARIH1 ([ARIH1 Products](#))

Background: E3 ubiquitin-protein ligase ARIH1 (EC 2.3.2.31) (H7-AP2) (HHARI) (Monocyte protein 6) (MOP-6) (Protein ariadne-1 homolog) (ARI-1) (UbcH7-binding protein) (UbcM4-interacting protein) (Ubiquitin-conjugating enzyme E2-binding protein 1),FUNCTION: E3 ubiquitin-protein ligase, which catalyzes ubiquitination of target proteins together with ubiquitin-conjugating enzyme E2 UBE2L3 (PubMed:15236971, PubMed:21532592, PubMed:24076655, PubMed:27565346, PubMed:23707686). Acts as an atypical E3 ubiquitin-protein ligase by working together with cullin-RING ubiquitin ligase (CRL) complexes and initiating ubiquitination of CRL substrates: associates with CRL complexes and specifically mediates addition of the first ubiquitin on CRLs targets (PubMed:27565346). The initial ubiquitin is then elongated by CDC34/UBE2R1 and UBE2R2 (PubMed:27565346). E3 ubiquitin-protein ligase activity is activated upon binding to neddylated cullin-RING ubiquitin ligase complexes (PubMed:24076655, PubMed:27565346). Plays a role in protein translation in response to DNA damage by mediating ubiquitination of EIF4E2, the consequences of EIF4E2 ubiquitination are however unclear (PubMed:25624349). According to a report, EIF4E2 ubiquitination leads to promote EIF4E2 cap-binding and protein translation arrest (PubMed:25624349). According to another report EIF4E2 ubiquitination leads to its subsequent degradation (PubMed:14623119). Acts as the ligase involved in ISGylation of EIF4E2 (PubMed:17289916). In vitro, controls the degradation of the LINC (LInker of Nucleoskeleton and Cytoskeleton) complex member SUN2 and may therefore have a role in the formation and localization of the LINC complex, and as a consequence, nuclear subcellular localization and nuclear morphology (PubMed:29689197). {ECO:0000269|PubMed:14623119, ECO:0000269|PubMed:15236971, ECO:0000269|PubMed:17289916, ECO:0000269|PubMed:21532592, ECO:0000269|PubMed:23707686, ECO:0000269|PubMed:24076655, ECO:0000269|PubMed:25624349, ECO:0000269|PubMed:27565346, ECO:0000269|PubMed:29689197}.

Target Details

Molecular Weight: 64.1 kDa

UniProt: [Q9Y4X5](#)

Application Details

Application Notes: In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.

Comment: ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from *Nicotiana tabacum* c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.

During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: The buffer composition is at the discretion of the manufacturer. If you have a special request, please contact us.

Handling Advice: Avoid repeated freeze-thaw cycles.

Storage: -80 °C

Storage Comment: Store at -80°C.

Expiry Date: Unlimited (if stored properly)



Image 1. „Crystallography Grade“ protein due to multi-step, protein-specific purification process