antibodies - online.com







ARL4C Protein (AA 2-192) (His tag)



Image



()	ve	K\ /		A .
	\cup	1 V/	Щ.	V۷

Overview		
Quantity:	1 mg	
Target:	ARL4C	
Protein Characteristics:	AA 2-192	
Origin:	Human	
Source:	Escherichia coli (E. coli)	
Protein Type:	Recombinant	
Purification tag / Conjugate:	This ARL4C protein is labelled with His tag.	
Application:	ELISA, Western Blotting (WB), Crystallization (Crys), SDS-PAGE (SDS)	
Product Details		
Sequence:	GNISSNISAF QSLHIVMLGL DSAGKTTVLY RLKFNEFVNT VPTIGFNTEK IKLSNGTAKG	

Application:	ELISA, Western Blotting (WB), Crystallization (Crys), SDS-PAGE (SDS)	
Product Details		
Sequence:	GNISSNISAF QSLHIVMLGL DSAGKTTVLY RLKFNEFVNT VPTIGFNTEK IKLSNGTAKG	
	ISCHFWDVGG QEKLRPLWKS YSRCTDGIIY VVDSVDVDRL EEAKTELHKV TKFAENQGTP	
	LLVIANKQDL PKSLPVAEIE KQLALHELIP ATTYHVQPAC AIIGEGLTEG MDKLYEMILK	
	RRKSLKQKKK R	
	Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a	
	special request, please contact us.	
Characteristics:	 Made in Germany - from design to production - by highly experienced protein experts. Human ARL4C Protein (raised in E. Coli) purified by multi-step, protein-specific process to 	
	ensure crystallization grade.	
	State-of-the-art algorithm used for plasmid design (Gene synthesis).	
	This protein is a made to order protein and will be made for the first time for your order. Our	
	experts in the lab will ensure that you receive a correctly folded protein.	

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).

When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered. The concentration of our recombinant proteins is measured using the absorbance at 280nm. The protein's absorbance will be measured in several dilutions and is measured against its

The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

Purification:

Two step purification of proteins expressed in bacterial culture:

- 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.
- 2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

Purity:

>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.

Sterility:

0.22 µm filtered

Endotoxin Level:

Endotoxin has not been removed. Please contact us if you require endotoxin removal.

Grade:

Crystallography grade

specific reference buffer.

Target Details

Target:	ARL4C
Alternative Name: ARL4C (ARL4C Products)	
Background: Small GTP-binding protein which cycles between an inactive GDP-bound and an active GDP-bound and active GDP-bound active GDP-bound and active GDP-bound Active GDP	
	bound form, and the rate of cycling is regulated by guanine nucleotide exchange factors (GEF) and GTPase-activating proteins (GAP). GTP-binding protein that does not act as an allosteric

activator of the cholera toxin catalytic subunit. May be involved in transport between a perinuclear compartment and the plasma membrane, apparently linked to the ABCA1-mediated cholesterol secretion pathway. Recruits CYTH1, CYTH2, CYTH3 and CYTH4 to the plasma membrane in the GDP-bound form. Regulates the microtubule-dependent intracellular vesicular transport from early endosome to recycling endosome process. {ECO:0000269|PubMed:15147902, ECO:0000269|PubMed:17398095, ECO:0000269|PubMed:19409876}.

Molecular Weight:

22.3 kDa Including tag.

UniProt:

P56559

Application Details

Application Notes: In addition to the applications listed above we expect the protein to work for fu	
	as well. As the protein has not been tested for functional studies yet we cannot offer a
	guarantee though.

Comment:

In cases in which it is highly likely that the recombinant protein with the default tag will be insoluble our protein lab may suggest a higher molecular weight tag (e.g. GST-tag) instead to increase solubility. We will discuss all possible options with you in detail to assure that you receive your protein of interest.

Restrictions:

For Research Use only

Handling

Format:	Liquid	
Buffer:	100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer.	
Handling Advice:	Avoid repeated freeze-thaw cycles.	
Storage:	-80 °C	
Storage Comment:	Store at -80°C.	
Expiry Date:	Unlimited (if stored properly)	



Image 1. "Crystallography Grade" protein due to multi-step, protein-specific purification process