antibodies .- online.com





ATG3 Protein (AA 1-314) (His tag)



Image



Go to Product page

Overview

| Quantity: | 1 mg |
|-------------------------------|--|
| Target: | ATG3 |
| Protein Characteristics: | AA 1-314 |
| Origin: | Human |
| Source: | Insect Cells |
| Protein Type: | Recombinant |
| Purification tag / Conjugate: | This ATG3 protein is labelled with His tag. |
| Application: | ELISA, SDS-PAGE (SDS), Western Blotting (WB), Crystallization (Crys) |

Product Details

Sequence:

MQNVINTVKG KALEVAEYLT PVLKESKFKE TGVITPEEFV AAGDHLVHHC PTWQWATGEE LKVKAYLPTG KQFLVTKNVP CYKRCKQMEY SDELEAIIEE DDGDGGWVDT YHNTGITGIT EAVKEITLEN KDNIRLQDCS ALCEEEEDED EGEAADMEEY EESGLLETDE ATLDTRKIVE ACKAKTDAGG EDAILQTRTY DLYITYDKYY QTPRLWLFGY DEQRQPLTVE HMYEDISQDH VKKTVTIENH PHLPPPPMCS VHPCRHAEVM KKIIETVAEG GGELGVHMYL LIFLKFVQAV

IPTIEYDYTR HFTM

Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a special request, please contact us.

Characteristics:

- Made in Germany from design to production by highly experienced protein experts.
- Human ATG3 Protein (raised in Insect Cells) purified by multi-step, protein-specific process to ensure crystallization grade.
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made to order protein and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).

When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered.

The concentration of our recombinant proteins is measured using the absorbance at 280nm.

The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.

The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

Purification:

Two step purification of proteins expressed in baculovirus infected SF9 insect cells:

- 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.
- 2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

Purity:

>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.

Sterility:

0.22 µm filtered

Endotoxin Level:

Protein is endotoxin free.

Grade:

Crystallography grade

Target Details

| Target: | ATG3 |
|-------------------|---|
| Alternative Name: | ATG3 (ATG3 Products) |
| Background: | E2 conjugating enzyme required for the cytoplasm to vacuole transport (Cvt), autophagy, and |

| mitochondrial homeostasis. Responsible for the E2-like covalent binding of | | |
|---|--|--|
| phosphatidylethanolamine to the C-terminal Gly of ATG8-like proteins (GABARAP, GABARAPL1, | | |
| GABARAPL2 or MAP1LC3A). The ATG12-ATG5 conjugate plays a role of an E3 and promotes | | |
| the transfer of ATG8-like proteins from ATG3 to phosphatidylethanolamine (PE). This step is | | |
| required for the membrane association of ATG8-like proteins. The formation of the ATG8- | | |
| phosphatidylethanolamine conjugates is essential for autophagy and for the cytoplasm to | | |
| vacuole transport (Cvt). Preferred substrate is MAP1LC3A. Also acts as an autocatalytic E2-like | | |
| enzyme, catalyzing the conjugation of ATG12 to itself, ATG12 conjugation to ATG3 playing a | | |
| role in mitochondrial homeostasis but not in autophagy. ATG7 (E1-like enzyme) facilitates this | | |
| reaction by forming an E1-E2 complex with ATG3. Promotes primary ciliogenesis by removing | | |
| OFD1 from centriolar satellites via the autophagic pathway. {ECO:0000269 PubMed:11825910, | | |
| ECO:0000269 PubMed:12207896, ECO:0000269 PubMed:12890687, | | |
| ECO:0000269 PubMed:16704426, ECO:0000269 PubMed:20723759}. | | |

| Molecular Weight: | 36.8 kDa Including tag. |
|-------------------|-------------------------|
| UniProt: | Q9NT62 |
| Pathways: | Autophagy |

Application Details

| Application Notes: | In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though. |
|--------------------|---|
| Comment: | In cases in which it is highly likely that the recombinant protein with the default tag will be insoluble our protein lab may suggest a higher molecular weight tag (e.g. GST-tag) instead to increase solubility. We will discuss all possible options with you in detail to assure that you receive your protein of interest. |
| Restrictions: | For Research Use only |

Handling

| Format: | Liquid |
|------------------|--|
| Buffer: | 100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer. |
| Handling Advice: | Avoid repeated freeze-thaw cycles. |
| Storage: | -80 °C |

Handling

| Storage Comment: | Store at -80°C. |
|------------------|--------------------------------|
| Expiry Date: | Unlimited (if stored properly) |

Images



Image 1. "Crystallography Grade" protein due to multi-step, protein-specific purification process