antibodies .- online.com





CACNA1G Protein (AA 1855-2377) (His tag)



Overview

Quantity:	1 mg	
Target:	CACNA1G	
Protein Characteristics:	AA 1855-2377	
Origin:	Human	
Source:	Insect Cells	
Protein Type:	Recombinant	
Purification tag / Conjugate:	This CACNA1G protein is labelled with His tag.	
Application:	Crystallization (Crys), ELISA, SDS-PAGE (SDS), Western Blotting (WB)	

Product Details

Sequence:

EESNKEAKEE AELEAELELE MKTLSPQPHS PLGSPFLWPG VEGPDSPDSP KPGALHPAAH ARSASHFSLE HPTDRQLFDT ISLLIQGSLE WELKLMDELA GPGGQPSAFP SAPSLGGSDP QIPLAEMEAL SLTSEIVSEP SCSLALTDDS LPDDMHTLLL SALESNMQPH PTELPGPDLL TVRKSGVSRT HSLPNDSYMC RHGSTAEGPL GHRGWGLPKA QSGSVLSVHS QPADTSYILQ LPKDAPHLLQ PHSAPTWGTI PKLPPPGRSP LAQRPLRRQA AIRTDSLDVQ GLGSREDLLA EVSGPSPPLA RAYSFWGQSS TQAQQHSRSH SKISKHMTPP APCPGPEPNW GKGPPETRSS LELDTELSWI SGDLLPPGGQ EEPPSPRDLK KCYSVEAQSC QRRPTSWLDE QRRHSIAVSC LDSGSQPHLG TDPSNLGGQP LGGPGSRPKK KLSPPSITID PPESQGPRTP PSPGICLRRR APSSDSKDPL ASGPPDSMAA SPSPKKDVLS LSGLSSDPAD LDP

Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a special request, please contact us.

Characteristics:

- Made in Germany from design to production by highly experienced protein experts.
- Human CACNA1G Protein (raised in Insect Cells) purified by multi-step, protein-specific process to ensure crystallization grade.
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made to order protein and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).

When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered.

The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.

The concentration of our recombinant proteins is measured using the absorbance at 280nm.

The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

Purification:

Two step purification of proteins expressed in baculovirus infected SF9 insect cells:

- 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.
- 2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

Purity: >95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.

Sterility: 0.22 µm filtered

Endotoxin Level: Protein is endotoxin free.

Grade: Crystallography grade

Target Details

Target:	CACNA1G		
Alternative Name:	CACNA1G (CACNA1G Products)		
Background:	Voltage-sensitive calcium channels (VSCC) mediate the entry of calcium ions into excitable		
	cells and are also involved in a variety of calcium-dependent processes, including muscle		
	contraction, hormone or neurotransmitter release, gene expression, cell motility, cell division		
	and cell death. The isoform alpha-1G gives rise to T-type calcium currents. T-type calcium		
	channels belong to the "low-voltage activated (LVA)" group and are strongly blocked by		
	mibefradil. A particularity of this type of channel is an opening at quite negative potentials and		
	voltage-dependent inactivation. T-type channels serve pacemaking functions in both central		
	neurons and cardiac nodal cells and support calcium signaling in secretory cells and vascular		
	smooth muscle. They may also be involved in the modulation of firing patterns of neurons		
	which is important for information processing as well as in cell growth processes.		
Molecular Weight:	56.6 kDa Including tag.		
UniProt:	043497		
Application Details			
Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies		
	as well. As the protein has not been tested for functional studies yet we cannot offer a gurantee		
	though.		
Comment:	In cases in which it is highly likely that the recombinant protein with the default tag will be		
	insoluble our protein lab may suggest a higher molecular weight tag (e.g. GST-tag) instead to		
	increase solubility. We will discuss all possible options with you in detail to assure that you		
	receive your protein of interest.		
Restrictions:	For Research Use only		
Handling			
Format:	Liquid		
Buffer:	100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer.		
Handling Advice:	Avoid repeated freeze-thaw cycles.		
Storage:	-80 °C		
Storage Comment:	Store at -80°C.		

1	1 11	١٠
\vdash	land	∥n∩
1	ıarıa	11110

Expiry Date:

Unlimited (if stored properly)