antibodies -online.com





CACNA1C Protein (AA 406-524) (His tag)



()	11/0	K\ /	iew	1
	\cup	'I V/I	$\square \vee \vee$	ı

Overview	
Quantity:	1 mg
Target:	CACNA1C
Protein Characteristics:	AA 406-524
Origin:	Human
Source:	Escherichia coli (E. coli)
Protein Type:	Recombinant
Purification tag / Conjugate:	This CACNA1C protein is labelled with His tag.
Application:	Western Blotting (WB), SDS-PAGE (SDS), ELISA, Crystallization (Crys)
Product Details	
Sequence:	GEFSKEREKA KARGDFQKLR EKQQLEEDLK GYLDWITQAE DIDPENEDEG MDEEKPRNMS
	MPTSETESVN TENVAGGDIE GENCGARLAH RISKSKFSRY WRRWNRFCRR KCRAAVKSN
	Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a
	special request, please contact us.
Characteristics:	 Made in Germany - from design to production - by highly experienced protein experts. Human CACNA1C Protein (raised in E. Coli) purified by multi-step, protein-specific process to ensure crystallization grade. State-of-the-art algorithm used for plasmid design (Gene synthesis).
	This protein is a made to order protein and will be made for the first time for your order. Our
	experts in the lab will ensure that you receive a correctly folded protein.
	The big advantage of ordering our made-to-order proteins in comparison to ordering custom
	made proteins from other companies is that there is no financial obligation in case the protein

cannot be expressed or purified.

In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).

When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered. The concentration of our recombinant proteins is measured using the absorbance at 280nm. The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.

The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

Purification:

Two step purification of proteins expressed in bacterial culture:

- 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.
- 2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

Purity:

>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.

Sterility:

0.22 µm filtered

Endotoxin Level:

Endotoxin has not been removed. Please contact us if you require endotoxin removal.

Grade:

Crystallography grade

Target Details

Target:	CACNA1C	
Alternative Name:	CACNA1C (CACNA1C Products)	
Background:	Voltage-sensitive calcium channels (VSCC) mediate the entry of calcium ions into excitable	
	cells and are also involved in a variety of calcium-dependent processes, including muscle	
	contraction, hormone or neurotransmitter release, gene expression, cell motility, cell division	
	and cell death. The isoform alpha-1C gives rise to L-type calcium currents. Long-lasting (L-type)	
	calcium channels belong to the 'high-voltage activated' (HVA) group. They are blocked by	

Expiry Date:

	dihydropyridines (DHP), phenylalkylamines, benzothiazepines, and by omega-agatoxin-IIIA	
	(omega-Aga-IIIA). They are however insensitive to omega-conotoxin-GVIA (omega-CTx-GVIA)	
	and omega-agatoxin-IVA (omega-Aga-IVA). Calcium channels containing the alpha-1C subunit	
	play an important role in excitation-contraction coupling in the heart. The various isoforms	
	display marked differences in the sensitivity to DHP compounds. Binding of calmodulin or	
	CABP1 at the same regulatory sites results in an opposit effects on the channel function.	
	{ECO:0000269 PubMed:12176756, ECO:0000269 PubMed:17071743,	
	ECO:0000269 PubMed:7737988, ECO:0000269 PubMed:8392192,	
	ECO:0000269 PubMed:9013606, ECO:0000269 PubMed:9607315}.	
Molecular Weight:	14.9 kDa Including tag.	
UniProt:	Q13936	
Pathways:	Hormone Transport, Carbohydrate Homeostasis	
Application Details		
Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies	
	as well. As the protein has not been tested for functional studies yet we cannot offer a gurantee though.	
Comment:	In cases in which it is highly likely that the recombinant protein with the default tag will be	
	insoluble our protein lab may suggest a higher molecular weight tag (e.g. GST-tag) instead to	
	increase solubility. We will discuss all possible options with you in detail to assure that you	
	receive your protein of interest.	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Buffer:	100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer.	
Handling Advice:	Avoid repeated freeze-thaw cycles.	
Storage:	-80 °C	
Storage Comment:	Store at -80°C.	

Unlimited (if stored properly)