# antibodies .- online.com





# CBLC Protein (AA 1-474) (His tag)



**Image** 



#### Overview

Quantity:	1 mg
Target:	CBLC
Protein Characteristics:	AA 1-474
Origin:	Human
Source:	Insect Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This CBLC protein is labelled with His tag.
Application:	ELISA, Western Blotting (WB), Crystallization (Crys), SDS-PAGE (SDS)

#### **Product Details**

### Sequence:

MALAVAPWGR QWEEARALGR AVRMLQRLEE QCVDPRLSVS PPSLRDLLPR TAQLLREVAH SRRAAGGGGP GGPGGSGDFL LIYLANLEAK SRQVAALLPP RGRRSANDEL FRAGSRLRRQ LAKLAIIFSH MHAELHALFP GGKYCGHMYQ LTKAPAHTFW RESCGARCVL PWAEFESLLG TCHPVEPGCT ALALRTTIDL TCSGHVSIFE FDVFTRLFQP WPTLLKNWQL LAVNHPGYMA FLTYDEVQER LQACRDKPGS YIFRPSCTRL GQWAIGYVSS DGSILQTIPA NKPLSQVLLE GQKDGFYLYP DGKTHNPDLT ELGQAEPQQR IHVSEEQLQL YWAMDSTFEL CKICAESNKD VKIEPCGHLL CSCCLAAWQH SDSQTCPFCR CEIKGWEAVS IYQFHGQATA EDSGNSSDQE GRELELGQVP LSAPPLPPRP DLPPRKPRNA QPKVRLLKGN SPPAALGPQD PAPA

Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a special request, please contact us.

#### Characteristics:

- Made in Germany from design to production by highly experienced protein experts.
- Human CBLC Protein (raised in Insect Cells) purified by multi-step, protein-specific process

to ensure crystallization grade.

• State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made to order protein and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).

When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered. The concentration of our recombinant proteins is measured using the absorbance at 280nm.

The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.

The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

## Purification:

Two step purification of proteins expressed in baculovirus infected SF9 insect cells:

- 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.
- Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

Purity:

>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.

Sterility:

0.22 µm filtered

Endotoxin Level:

Protein is endotoxin free.

Grade:

Crystallography grade

## **Target Details**

Target: CBLC

Alternative Name:

CBLC (CBLC Products)


# Target Details

ranger Betane	
Background:	Acts as an E3 ubiquitin-protein ligase, which accepts ubiquitin from specific E2 ubiquitin-
	conjugating enzymes, and then transfers it to substrates promoting their degradation by the
	proteasome. Functionally coupled with the E2 ubiquitin-protein ligases UB2D1, UB2D2 and
	UB2D3. Regulator of EGFR mediated signal transduction, upon EGF activation, ubiquitinates
	EGFR. Isoform 1, but not isoform 2, inhibits EGF stimulated MAPK1 activation. Promotes
	ubiquitination of SRC phosphorylated at 'Tyr-419'. In collaboration with CD2AP may act as
	regulatory checkpoint for Ret signaling by modulating the rate of RET degradation after ligand
	activation, CD2AP converts it from an inhibitor to a promoter of RET degradation, the function
	limits the potency of GDNF on neuronal survival. {ECO:0000269 PubMed:10362357,
	ECO:0000269 PubMed:14661060, ECO:0000269 PubMed:18753381,
	ECO:0000269 PubMed:20525694, ECO:0000269 PubMed:23145173}.
Molecular Weight:	53.4 kDa Including tag.
UniProt:	Q9ULV8
Pathways:	EGFR Signaling Pathway
Application Details	
Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies
	as well. As the protein has not been tested for functional studies yet we cannot offer a gurantee
	though.
Comment:	In cases in which it is highly likely that the recombinant protein with the default tag will be
	insoluble our protein lab may suggest a higher molecular weight tag (e.g. GST-tag) instead to
	increase solubility. We will discuss all possible options with you in detail to assure that you
	receive your protein of interest.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.

Expiry Date:

Unlimited (if stored properly)

Images



**Image 1.** "Crystallography Grade" protein due to multi-step, protein-specific purification process