

Datasheet for ABIN3091242 CD5L Protein (AA 20-347) (His tag)



[Go to Product page](#)

1 Image

Overview

Quantity:	1 mg
Target:	CD5L
Protein Characteristics:	AA 20-347
Origin:	Human
Source:	Insect Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This CD5L protein is labelled with His tag.
Application:	Western Blotting (WB), SDS-PAGE (SDS), ELISA, Crystallization (Crys)

Product Details

Sequence:	<p>SPSGVRLVGG LHRCEGRVEV EQKGQWGTV C DDGWDIKDVA VLCRELGCGA ASGTPSGILY EPPAEKEQKV LIQSVSCTGT EDTLAQCEQE EVYDCSHDED AGASCENPES SFSPVPEGVR LADGPGHCKG RVEVKHQNQW YTV CQTGWSL RAAKVVC RQL GCGRAVL TQK RCNKHAYGRK PIWLSQMSCS GREATLQDCP SGPWGKNTCN HEDT WVECE DPFDLRLVGG DNLCSGRLEV LHKG VWG SVC DDNWGEKEDQ VVCKQLGCGK SLSPSFRDRK CYGPGVGRIW LDNVRCSGEE QSLEQCQHRF WGFHDCTHQE DVAVICSG</p> <p>Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a special request, please contact us.</p>
Characteristics:	<ul style="list-style-type: none"> • Made in Germany - from design to production - by highly experienced protein experts. • Human CD5L Protein (raised in Insect Cells) purified by multi-step, protein-specific process to ensure crystallization grade. • State-of-the-art algorithm used for plasmid design (Gene synthesis).

Product Details

This protein is a made to order protein and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).

When you order this made-to-order protein you will only pay upon receipt of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered.

The concentration of our recombinant proteins is measured using the absorbance at 280nm. The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.

The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

Purification:	Two step purification of proteins expressed in baculovirus infected SF9 insect cells: <ol style="list-style-type: none">1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.
---------------	--

Purity:	>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.
---------	--

Sterility:	0.22 µm filtered
------------	------------------

Endotoxin Level:	Protein is endotoxin free.
------------------	----------------------------

Grade:	Crystallography grade
--------	-----------------------

Target Details

Target:	CD5L
---------	------

Alternative Name:	CD5L (CD5L Products)
-------------------	--

Background:	Secreted protein that acts as a key regulator of lipid synthesis: mainly expressed by
-------------	---

Target Details

macrophages in lymphoid and inflamed tissues and regulates mechanisms in inflammatory responses, such as infection or atherosclerosis. Able to inhibit lipid droplet size in adipocytes. Following incorporation into mature adipocytes via CD36-mediated endocytosis, associates with cytosolic FASN, inhibiting fatty acid synthase activity and leading to lipolysis, the degradation of triacylglycerols into glycerol and free fatty acids (FFA). CD5L-induced lipolysis occurs with progression of obesity: participates to obesity-associated inflammation following recruitment of inflammatory macrophages into adipose tissues, a cause of insulin resistance and obesity-related metabolic disease. Regulation of intracellular lipids mediated by CD5L has a direct effect on transcription regulation mediated by nuclear receptors ROR-gamma (RORC). Acts as a key regulator of metabolic switch in T-helper Th17 cells. Regulates the expression of pro-inflammatory genes in Th17 cells by altering the lipid content and limiting synthesis of cholesterol ligand of RORC, the master transcription factor of Th17-cell differentiation. CD5L is mainly present in non-pathogenic Th17 cells, where it decreases the content of polyunsaturated fatty acyls (PUFA), affecting two metabolic proteins MSMO1 and CYP51A1, which synthesize ligands of RORC, limiting RORC activity and expression of pro-inflammatory genes. Participates in obesity-associated autoimmunity via its association with IgM, interfering with the binding of IgM to Fc α /mu receptor and enhancing the development of long-lived plasma cells that produce high-affinity IgG autoantibodies (By similarity). Also acts as an inhibitor of apoptosis in macrophages: promotes macrophage survival from the apoptotic effects of oxidized lipids in case of atherosclerosis (PubMed:24295828). Involved in early response to microbial infection against various pathogens by acting as a pattern recognition receptor and by promoting autophagy (PubMed:16030018, PubMed:24223991, PubMed:24583716, PubMed:25713983). {ECO:0000250|UniProtKB:Q9QWK4, ECO:0000269|PubMed:16030018, ECO:0000269|PubMed:24223991, ECO:0000269|PubMed:24295828, ECO:0000269|PubMed:24583716, ECO:0000269|PubMed:25713983}.

Molecular Weight:	37.0 kDa Including tag.
-------------------	-------------------------

UniProt:	O43866
----------	------------------------

Application Details

Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.
--------------------	--

Comment:	In cases in which it is highly likely that the recombinant protein with the default tag will be insoluble our protein lab may suggest a higher molecular weight tag (e.g. GST-tag) instead to
----------	---

Application Details

increase solubility. We will discuss all possible options with you in detail to assure that you receive your protein of interest.

Restrictions: For Research Use only

Handling

Format:	Liquid
Buffer:	100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	Unlimited (if stored properly)

Images



Image 1. „Crystallography Grade“ protein due to multi-step, protein-specific purification process