

Datasheet for ABIN3091775 **C4A Protein (AA 20-675) (His tag)**



Overview

Quantity:	1 mg
Target:	C4A (C4a)
Protein Characteristics:	AA 20-675
Origin:	Human
Source:	Insect Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This C4A protein is labelled with His tag.
Application:	Western Blotting (WB), SDS-PAGE (SDS), ELISA, Crystallization (Crys)

Product Details

Sequence:

KPRLLLFSPS VVHLGVPLSV GVQLQDVPRG QVVKGSVFLR NPSRNNVPCS PKVDFTLSSE RDFALLSLQV PLKDAKSCGL HQLLRGPEVQ LVAHSPWLKD SLSRTTNIQG INLLFSSRRG HLFLQTDQPI YNPGQRVRYR VFALDQKMRP STDTITVMVE NSHGLRVRKK EVYMPSSIFQ DDFVIPDISE PGTWKISARF SDGLESNSST QFEVKKYVLP NFEVKITPGK PYILTVPGHL DEMQLDIQAR YIYGKPVQGV AYVRFGLLDE DGKKTFFRGL ESQTKLVNGQ SHISLSKAEF QDALEKLNMG ITDLQGLRLY VAAAIIESPG GEMEEAELTS WYFVSSPFSL DLSKTKRHLV PGAPFLLQAL VREMSGSPAS GIPVKVSATV SSPGSVPEVQ DIQQNTDGSG QVSIPIIIPQ TISELQLSVS AGSPHPAIAR LTVAAPPSGG PGFLSIERPD SRPPRVGDTL NLNLRAVGSG ATFSHYYYMI LSRGQIVFMN REPKRTLTSV SVFVDHHLAP SFYFVAFYYH GDHPVANSLR VDVQAGACEG KLELSVDGAK QYRNGESVKL HLETDSLALV ALGALDTALY AAGSKSHKPL NMGKVFEAMN SYDLGCGPGG GDSALQVFQA AGLAFSDGDQ WTLSRKRLSC PKEKTT

Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a

Product Details special request, please contact us. Characteristics: · Made in Germany - from design to production - by highly experienced protein experts. · Human C4A Protein (raised in Insect Cells) purified by multi-step, protein-specific process to ensure crystallization grade. • State-of-the-art algorithm used for plasmid design (Gene synthesis). This protein is a made to order protein and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein. The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified. In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization). When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered. The concentration of our recombinant proteins is measured using the absorbance at 280nm. The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer. The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein. Purification: Two step purification of proteins expressed in baculovirus infected SF9 insect cells: 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE. 2. Protein containing fractions of the best purification are subjected to second purification step

through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot. Purity: 0.22 µm filtered Sterility: Endotoxin Level: Protein is endotoxin free. Grade: Crystallography grade

Target Details

Target:	C4A (C4a)	
Alternative Name:	C4A (C4a Products)	
Background:	Non-enzymatic component of C3 and C5 convertases and thus essential for the propagation of	
	the classical complement pathway. Covalently binds to immunoglobulins and immune	
	complexes and enhances the solubilization of immune aggregates and the clearance of IC	
	through CR1 on erythrocytes. C4A isotype is responsible for effective binding to form amide	
	bonds with immune aggregates or protein antigens, while C4B isotype catalyzes the	
	transacylation of the thioester carbonyl group to form ester bonds with carbohydrate antigens.	
	Derived from proteolytic degradation of complement C4, C4a anaphylatoxin is a mediator of	
	local inflammatory process. It induces the contraction of smooth muscle, increases vascular	
	permeability and causes histamine release from mast cells and basophilic leukocytes.	
Molecular Weight:	72.6 kDa Including tag.	
UniProt:	P0C0L4	
Pathways:	Complement System	
Application Details		
Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies	
	as well. As the protein has not been tested for functional studies yet we cannot offer a gurantee	
	though.	
Comment:	In cases in which it is highly likely that the recombinant protein with the default tag will be	
	insoluble our protein lab may suggest a higher molecular weight tag (e.g. GST-tag) instead to	
	increase solubility. We will discuss all possible options with you in detail to assure that you	
	receive your protein of interest.	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Buffer:	100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer.	
Handling Advice:	Avoid repeated freeze-thaw cycles.	
Storage:	-80 °C	
Storage Comment:	Store at -80°C.	

\vdash	land	linc
	iaria	11116

Expiry Date:

Unlimited (if stored properly)