antibodies .- online.com





C5 Protein (AA 19-673) (His tag)



Go to Product pag

Overview

Quantity:	1 mg	
Target:	C5	
Protein Characteristics:	AA 19-673	
Origin:	Human	
Source:	Insect Cells	
Protein Type:	Recombinant	
Purification tag / Conjugate:	This C5 protein is labelled with His tag.	
Application:	Western Blotting (WB), SDS-PAGE (SDS), ELISA, Crystallization (Crys)	

Product Details

Sequence:

QEQTYVISAP KIFRVGASEN IVIQVYGYTE AFDATISIKS YPDKKFSYSS GHVHLSSENK
FQNSAILTIQ PKQLPGGQNP VSYVYLEVVS KHFSKSKRMP ITYDNGFLFI HTDKPVYTPD
QSVKVRVYSL NDDLKPAKRE TVLTFIDPEG SEVDMVEEID HIGIISFPDF KIPSNPRYGM
WTIKAKYKED FSTTGTAYFE VKEYVLPHFS VSIEPEYNFI GYKNFKNFEI TIKARYFYNK
VVTEADVYIT FGIREDLKDD QKEMMQTAMQ NTMLINGIAQ VTFDSETAVK ELSYYSLEDL
NNKYLYIAVT VIESTGGFSE EAEIPGIKYV LSPYKLNLVA TPLFLKPGIP YPIKVQVKDS
LDQLVGGVPV TLNAQTIDVN QETSDLDPSK SVTRVDDGVA SFVLNLPSGV TVLEFNVKTD
APDLPEENQA REGYRAIAYS SLSQSYLYID WTDNHKALLV GEHLNIIVTP KSPYIDKITH
YNYLILSKGK IIHFGTREKF SDASYQSINI PVTQNMVPSS RLLVYYIVTG EQTAELVSDS
VWLNIEEKCG NQLQVHLSPD ADAYSPGQTV SLNMATGMDS WVALAAVDSA VYGVQRGAKK
PLERVFQFLE KSDLGCGAGG GLNNANVFHL AGLTFLTNAN ADDSQENDEP CKEIL

Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a

special request, please contact us.

Characteristics:

- · Made in Germany from design to production by highly experienced protein experts.
- Human C5 Protein (raised in E. Coli) purified by multi-step, protein-specific process to ensure crystallization grade.
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made to order protein and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).

When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered.

The concentration of our recombinant proteins is measured using the absorbance at 280nm.

The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.

The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

Purification:

Two step purification of proteins expressed in bacterial culture:

- 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.
- 2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

Purity:

>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.

Sterility:

0.22 µm filtered

Endotoxin Level:

Protein is endotoxin free.

Grade:

Crystallography grade

Target Details

Target:	C5	
Alternative Name:	C5 (C5 Products)	
Background:	Activation of C5 by a C5 convertase initiates the spontaneous assembly of the late complement components, C5-C9, into the membrane attack complex. C5b has a transient binding site for C6. The C5b-C6 complex is the foundation upon which the lytic complex is assembled., Derived from proteolytic degradation of complement C5, C5 anaphylatoxin is a mediator of local inflammatory process. Binding to the receptor C5AR1 induces a variety of responses including intracellular calcium release, contraction of smooth muscle, increased vascular permeability, and histamine release from mast cells and basophilic leukocytes (PubMed:8182049). C5a is also a potent chemokine which stimulates the locomotion of polymorphonuclear leukocytes	
	and directs their migration toward sites of inflammation. {ECO:0000269 PubMed:8182049}.	
Molecular Weight:	74.2 kDa Including tag.	
UniProt:	P01031	
Pathways:	Complement System, Carbohydrate Homeostasis	
Application Details		
Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a gurantee though.	
Comment:	In cases in which it is highly likely that the recombinant protein with the default tag will be insoluble our protein lab may suggest a higher molecular weight tag (e.g. GST-tag) instead to increase solubility. We will discuss all possible options with you in detail to assure that you receive your protein of interest.	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Buffer:	100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer.	
Handling Advice:	Avoid repeated freeze-thaw cycles.	
Storage:	-80 °C	
Storage Comment:	Store at -80°C.	

1	1 11	١٠
\vdash	land	∥n∩
1	ıarıa	11110

Expiry Date:

Unlimited (if stored properly)