antibodies .- online.com



Datasheet for ABIN3091840

C-Type Lectin Domain Family 4, Member M (CLEC4M) (AA 71-399) protein (His tag)



Go to Product page



Overview

| Quantity: | 1 mg |
|-------------------------------|--|
| Target: | C-Type Lectin Domain Family 4, Member M (CLEC4M) |
| Protein Characteristics: | AA 71-399 |
| Origin: | Human |
| Source: | Insect Cells |
| Protein Type: | Recombinant |
| Purification tag / Conjugate: | His tag |
| Application: | Western Blotting (WB), SDS-PAGE (SDS), ELISA, Crystallization (Crys) |

Product Details

Sequence:

QVSKVPSSLS QEQSEQDAIY QNLTQLKAAV GELSEKSKLQ EIYQELTQLK AAVGELPEKS
KLQEIYQELT RLKAAVGELP EKSKLQEIYQ ELTRLKAAVG ELPEKSKLQE IYQELTRLKA
AVGELPEKSK LQEIYQELTE LKAAVGELPE KSKLQEIYQE LTQLKAAVGE LPDQSKQQQI
YQELTDLKTA FERLCRHCPK DWTFFQGNCY FMSNSQRNWH DSVTACQEVR AQLVVIKTAE
EQNFLQLQTS RSNRFSWMGL SDLNQEGTWQ WVDGSPLSPS FQRYWNSGEP NNSGNEDCAE
FSGSGWNDNR CDVDNYWICK KPAACFRDE

Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a special request, please contact us.

Characteristics:

- Made in Germany from design to production by highly experienced protein experts.
- Human CLEC4M Protein (raised in Insect Cells) purified by multi-step, protein-specific process to ensure crystallization grade.
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made to order protein and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).

When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered.

The concentration of our recombinant proteins is measured using the absorbance at 280nm.

The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.

The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

Purification:

Two step purification of proteins expressed in baculovirus infected SF9 insect cells:

- 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.
- 2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

Purity:

>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.

Sterility:

0.22 µm filtered

Endotoxin Level:

Protein is endotoxin free.

Grade:

Crystallography grade

Target Details

| Target: | C-Type Lectin Domain Family 4, Member M (CLEC4M) |
|-------------------|---|
| Alternative Name: | CLEC4M (CLEC4M Products) |
| Background: | Probable pathogen-recognition receptor involved in peripheral immune surveillance in liver. May |

mediate the endocytosis of pathogens which are subsequently degraded in lysosomal compartments. Is a receptor for ICAM3, probably by binding to mannose-like carbohydrates. {ECO:0000269|PubMed:11257134}., (Microbial infection) Acts as an attachment receptor for ebola virus (PubMed:12050398, PubMed:12502850). Acts as an attachment receptor for hepatitis C virus (PubMed:15371595, PubMed:16816373). Acts as an attachment receptor for HIV-1 (PubMed:21277928, PubMed:12502850). Acts as an attachment receptor for human coronavirus 229E (PubMed:17037540). Acts as an attachment receptor for human cytomegalovirus/HHV-5 (PubMed:12433371). Acts as an attachment receptor for influenzavirus (PubMed:21191006). Acts as an attachment receptor for SARS coronavirus (PubMed:15479853). Acts as an attachment receptor for west-nile virus (PubMed:16415006). Acts as an attachment receptor for Japanese encephalitis virus (PubMed:24623090). Acts as an attachment receptor for Marburg virus glycoprotein (PubMed:15479853). {ECO:0000269|PubMed:12050398, ECO:0000269|PubMed:12433371, ECO:0000269|PubMed:12502850, ECO:0000269|PubMed:15371595, ECO:0000269|PubMed:15479853, ECO:0000269|PubMed:16415006, ECO:0000269|PubMed:16816373, ECO:0000269|PubMed:17037540,

ECO:0000269|PubMed:21191006, ECO:0000269|PubMed:21277928,

Molecular Weight:

38.7 kDa Including tag.

ECO:0000269|PubMed:24623090}.

UniProt:

Q9H2X3

Application Details

Application Notes:

In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a gurantee though.

Comment:

In cases in which it is highly likely that the recombinant protein with the default tag will be insoluble our protein lab may suggest a higher molecular weight tag (e.g. GST-tag) instead to increase solubility. We will discuss all possible options with you in detail to assure that you receive your protein of interest.

Restrictions:

For Research Use only

Handling

Format:

Liquid

Handling

| Buffer: | 100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer. |
|------------------|--|
| Handling Advice: | Avoid repeated freeze-thaw cycles. |
| Storage: | -80 °C |
| Storage Comment: | Store at -80°C. |
| Expiry Date: | Unlimited (if stored properly) |

Images




Image 1. "Crystallography Grade" protein due to multi-step, protein-specific purification process