antibodies .- online.com





CYP1A2 Protein (AA 2-515) (His tag)





Go to Product page

Overview

Quantity:	1 mg
Target:	CYP1A2
Protein Characteristics:	AA 2-515
Origin:	Human
Source:	Insect Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This CYP1A2 protein is labelled with His tag.
Application:	Western Blotting (WB), SDS-PAGE (SDS), ELISA, Crystallization (Crys)

Product Details

Sequence:

ALSQSVPFSA TELLLASAIF CLVFWVLKGL RPRVPKGLKS PPEPWGWPLL GHVLTLGKNP HLALSRMSQR YGDVLQIRIG STPVLVLSRL DTIRQALVRQ GDDFKGRPDL YTSTLITDGQ SLTFSTDSGP VWAARRRLAQ NALNTFSIAS DPASSSSCYL EEHVSKEAKA LISRLQELMA GPGHFDPYNQ VVVSVANVIG AMCFGQHFPE SSDEMLSLVK NTHEFVETAS SGNPLDFFPI LRYLPNPALQ RFKAFNQRFL WFLQKTVQEH YQDFDKNSVR DITGALFKHS KKGPRASGNL IPQEKIVNLV NDIFGAGFDT VTTAISWSLM YLVTKPEIQR KIQKELDTVI GRERRPRLSD RPQLPYLEAF ILETFRHSSF LPFTIPHSTT RDTTLNGFYI PKKCCVFVNQ WQVNHDPELW EDPSEFRPER FLTADGTAIN KPLSEKMMLF GMGKRRCIGE VLAKWEIFLF LAILLQQLEF SVPPGVKVDL TPIYGLTMKH ARCEHVQARR FSIN

Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a special request, please contact us.

Characteristics:

- Made in Germany from design to production by highly experienced protein experts.
- Human CYP1A2 Protein (raised in Insect Cells) purified by multi-step, protein-specific process to ensure crystallization grade.
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made to order protein and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).

When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered.

The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.

The concentration of our recombinant proteins is measured using the absorbance at 280nm.

The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

Purification:

Two step purification of proteins expressed in baculovirus infected SF9 insect cells:

- 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.
- 2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

Purity: >95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.

Sterility: 0.22 µm filtered

Endotoxin Level: Protein is endotoxin free.

Grade: Crystallography grade

Target Details

Target:	CYP1A2
Alternative Name:	CYP1A2 (CYP1A2 Products)
Background:	Cytochromes P450 are a group of heme-thiolate monooxygenases. In liver microsomes, this enzyme is involved in an NADPH-dependent electron transport pathway. It oxidizes a variety of structurally unrelated compounds, including steroids, fatty acids, and xenobiotics. Most active in catalyzing 2-hydroxylation. Caffeine is metabolized primarily by cytochrome CYP1A2 in the liver through an initial N3-demethylation. Also acts in the metabolism of aflatoxin B1 and acetaminophen. Participates in the bioactivation of carcinogenic aromatic and heterocyclic amines. Catalizes the N-hydroxylation of heterocyclic amines and the O-deethylation of phenacetin. {ECO:0000269 PubMed:14725854}.
Molecular Weight:	59.1 kDa Including tag.
UniProt:	P05177
Application Details	
Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a gurantee though.
Comment:	In cases in which it is highly likely that the recombinant protein with the default tag will be insoluble our protein lab may suggest a higher molecular weight tag (e.g. GST-tag) instead to increase solubility. We will discuss all possible options with you in detail to assure that you receive your protein of interest.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	Unlimited (if stored properly)



Image 1. "Crystallography Grade" protein due to multi-step, protein-specific purification process