# antibodies .- online.com





# DDX21 Protein (AA 1-783) (Strep Tag)



Go to Product page

### Overview

Quantity:	1 mg
Target:	DDX21
Protein Characteristics:	AA 1-783
Origin:	Human
Source:	Tobacco (Nicotiana tabacum)
Protein Type:	Recombinant
Purification tag / Conjugate:	This DDX21 protein is labelled with Strep Tag.
Application:	ELISA, Western Blotting (WB), SDS-PAGE (SDS)

# **Product Details**

Sequence:

MPGKLRSDAG LESDTAMKKG ETLRKQTEEK EKKEKPKSDK TEEIAEEEET VFPKAKQVKK KAEPSEVDMN SPKSKKAKKK EEPSQNDISP KTKSLRKKKE PIEKKVVSSK TKKVTKNEEP SEEEIDAPKP KKMKKEKEMN GETREKSPKL KNGFPHPEPD CNPSEAASEE SNSEIEQEIP VEQKEGAFSN FPISEETIKL LKGRGVTFLF PIQAKTFHHV YSGKDLIAQA RTGTGKTFSF AIPLIEKLHG ELQDRKRGRA PQVLVLAPTR ELANQVSKDF SDITKKLSVA CFYGGTPYGG QFERMRNGID ILVGTPGRIK DHIQNGKLDL TKLKHVVLDE VDQMLDMGFA DQVEEILSVA YKKDSEDNPQ TLLFSATCPH WVFNVAKKYM KSTYEQVDLI GKKTQKTAIT VEHLAIKCHW TQRAAVIGDV IRVYSGHQGR TIIFCETKKE AQELSQNSAI KQDAQSLHGD IPQKQREITL KGFRNGSFGV LVATNVAARG LDIPEVDLVI QSSPPKDVES YIHRSGRTGR AGRTGVCICF YQHKEEYQLV QVEQKAGIKF KRIGVPSATE IIKASSKDAI RLLDSVPPTA ISHFKQSAEK LIEEKGAVEA LAAALAHISG ATSVDQRSLI NSNVGFVTMI LQCSIEMPNI SYAWKELKEQ LGEEIDSKVK GMVFLKGKLG VCFDVPTASV TEIQEKWHDS RRWQLSVATE QPELEGPREG

YGGFRGQREG SRGFRGQRDG NRRFRGQREG SRGPRGQRSG GGNKSNRSQN KGQKRSFSKA FGQ

Sequence without tag. The proposed Strep-Tag is based on experience s with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.

#### Characteristics:

# Key Benefits:

- Made in Germany from design to production by highly experienced protein experts.
- Protein expressed with ALiCE® and purified by multi-step, protein-specific process to ensure correct folding and modification.
- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

#### Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require posttranslational modifications.
- During lysate production, the cell wall and other cellular components that are not required for
  protein production are removed, leaving only the protein production machinery and the
  mitochondria to drive the reaction. During our lysate completion steps, the additional
  components needed for protein production (amino acids, cofactors, etc.) are added to
  produce something that functions like a cell, but without the constraints of a living system all that's needed is the DNA that codes for the desired protein!

## Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.
- We use the Expasy's ProtParam tool to determine the absorption coefficient of each protein.

# **Product Details**

#### Purification:

Two step purification of proteins expressed in Almost Living Cell-Free Expression System (ALiCE®):

- 1. In a first purification step, the protein is purified from the cleared cell lysate using StrepTag capture material. Eluate fractions are analyzed by SDS-PAGE.
- Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

Purity:

>80 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.

Endotoxin Level:

Low Endotoxin less than 1 EU/mg (< 0.1 ng/mg)

# **Target Details**

Target:

DDX21

Alternative Name:

DDX21 (DDX21 Products)

Background:

Nucleolar RNA helicase 2 (EC 3.6.4.13) (DEAD box protein 21) (Gu-alpha) (Nucleolar RNA helicase Gu) (Nucleolar RNA helicase II) (RH II/Gu), FUNCTION: RNA helicase that acts as a sensor of the transcriptional status of both RNA polymerase (Pol) I and II: promotes ribosomal RNA (rRNA) processing and transcription from polymerase II (Pol II) (PubMed:25470060, PubMed:28790157). Binds various RNAs, such as rRNAs, snoRNAs, 7SK and, at lower extent, mRNAs (PubMed:25470060). In the nucleolus, localizes to rDNA locus, where it directly binds rRNAs and snoRNAs, and promotes rRNA transcription, processing and modification. Required for rRNA 2'-O-methylation, possibly by promoting the recruitment of late-acting snoRNAs SNORD56 and SNORD58 with pre-ribosomal complexes (PubMed:25470060, PubMed:25477391). In the nucleoplasm, binds 7SK RNA and is recruited to the promoters of Pol II-transcribed genes: acts by facilitating the release of P-TEFb from inhibitory 7SK snRNP in a manner that is dependent on its helicase activity, thereby promoting transcription of its target genes (PubMed:25470060). Functions as a cofactor for JUN-activated transcription: required for phosphorylation of JUN at 'Ser-77' (PubMed:11823437, PubMed:25260534). Can unwind double-stranded RNA (helicase) and can fold or introduce a secondary structure to a singlestranded RNA (foldase) (PubMed:9461305). Together with SIRT7, required to prevent R-loopassociated DNA damage and transcription-associated genomic instability: deacetylation by SIRT7 activates the helicase activity, thereby overcoming R-loop-mediated stalling of RNA polymerases (PubMed:28790157). Involved in rRNA processing (PubMed:14559904, PubMed:18180292). May bind to specific miRNA hairpins (PubMed:28431233). Component of a multi-helicase-TICAM1 complex that acts as a cytoplasmic sensor of viral double-stranded

rarget betano	
	RNA (dsRNA) and plays a role in the activation of a cascade of antiviral responses including the
	induction of pro-inflammatory cytokines via the adapter molecule TICAM1 (By similarity).
	{ECO:0000250 UniProtKB:Q9JIK5, ECO:0000269 PubMed:11823437,
	ECO:0000269 PubMed:14559904, ECO:0000269 PubMed:18180292,
	ECO:0000269 PubMed:25260534, ECO:0000269 PubMed:25470060,
	ECO:0000269 PubMed:25477391, ECO:0000269 PubMed:28431233,
	ECO:0000269 PubMed:28790157, ECO:0000269 PubMed:9461305}.
Molecular Weight:	87.3 kDa
UniProt:	Q9NR30
Pathways:	SARS-CoV-2 Protein Interactome
Application Details	
Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies
	as well. As the protein has not been tested for functional studies yet we cannot offer a
	guarantee though.
Comment:	ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from
	Nicotiana tabacum c.v This contains all the protein expression machinery needed to produce
	even the most difficult-to-express proteins, including those that require post-translational
	modifications.
	During lysate production, the cell wall and other cellular components that are not required for
	protein production are removed, leaving only the protein production machinery and the
	mitochondria to drive the reaction. During our lysate completion steps, the additional
	components needed for protein production (amino acids, cofactors, etc.) are added to produce
	something that functions like a cell, but without the constraints of a living system - all that's
	needed is the DNA that codes for the desired protein!
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer. If you have a special request, please contact us.
Handling Advice:	Avoid repeated freeze-thaw cycles.

# Handling

Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	Unlimited (if stored properly)