
antibodies .- online.com





DHX16 Protein (AA 1-1041) (Strep Tag)



Image



Overview

Quantity:	1 mg
Target:	DHX16
Protein Characteristics:	AA 1-1041
Origin:	Human
Source:	Tobacco (Nicotiana tabacum)
Protein Type:	Recombinant
Purification tag / Conjugate:	This DHX16 protein is labelled with Strep Tag.
Application:	ELISA, Western Blotting (WB), SDS-PAGE (SDS)

Product Details

Sequence:

MATPAGLERW VQDELHSVLG LSERHVAQFL IGTAQRCTSA EEFVQRLRDT DTLDLSGPAR DFALRLWNKV PRKAVVEKPA RAAEREARAL LEKNRSYRLL EDSEESSEET VSRAGSSLQK KRKKRKHLRK KREEEEEEA SEKGKKKTGG SKQQTEKPES EDEWERTERE RLQDLEERDA FAERVRQRDK DRTRNVLERS DKKAYEEAQK RLKMAEEDRK AMVPELRKKS RREYLAKRER EKLEDLEAEL ADEEFLFGDV ELSRHERQEL KYKRRVRDLA REYRAAGEQE KLEATNRYHM PKETRGQPAR AVDLVEEESG APGEEQRRWE EARLGAASLK FGARDAASQE PKYQLVLEEE ETIEFVRATQ LQGDEEPSAP PTSTQAQQKE SIQAVRRSLP VFPFREELLA AIANHQVLII EGETGSGKTT QIPQYLFEEG YTNKGMKIAC TQPRRVAAMS VAARVAREMG VKLGNEVGYS IRFEDCTSER TVLRYMTDGM LLREFLSEPD LASYSVVMVD EAHERTLHTD ILFGLIKDVA RFRPELKVLV ASATMDTARF STFFDDAPVF RIPGRRFPVD IFYTKAPEAD YLEACVVSVL QIHVTQPPGD ILVFLTGQEE IEAACEMLQD RCRRLGSKIR ELLVLPIYAN LPSDMQARIF QPTPPGARKV VVATNIAETS LTIEGIIYVL DPGFCKQKSY NPRTGMESLT VTPCSKASAN

QRAGRAGRVA AGKCFRLYTA WAYQHELEET TVPEIQRTSL GNVVLLLKSL GIHDLMHFDF
LDPPPYETLL LALEQLYALG ALNHLGELTT SGRKMAELPV DPMLSKMILA SEKYSCSEEI
LTVAAMLSVN NSIFYRPKDK VVHADNARVN FFLPGGDHLV LLNVYTQWAE SGYSSQWCYE
NFVQFRSMRR ARDVREQLEG LLERVEVGLS SCQGDYIRVR KAITAGYFYH TARLTRSGYR
TVKQQQTVFI HPNSSLFEQQ PRWLLYHELV LTTKEFMRQV LEIESSWLLE VAPHYYKAKE
LEDPHAKKMP KKIGKTREEL G

Sequence without tag. The proposed Strep-Tag is based on experience s with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.

Characteristics:

Key Benefits:

- Made in Germany from design to production by highly experienced protein experts.
- Protein expressed with ALiCE® and purified by multi-step, protein-specific process to ensure correct folding and modification.
- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require posttranslational modifications.
- During lysate production, the cell wall and other cellular components that are not required for
 protein production are removed, leaving only the protein production machinery and the
 mitochondria to drive the reaction. During our lysate completion steps, the additional
 components needed for protein production (amino acids, cofactors, etc.) are added to
 produce something that functions like a cell, but without the constraints of a living system all that's needed is the DNA that codes for the desired protein!

Concentration:

The concentration of our recombinant proteins is measured using the absorbance at 280nm.

- The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.
- We use the Expasy's ProtParam tool to determine the absorption coefficient of each protein.

Purification:

Two step purification of proteins expressed in Almost Living Cell-Free Expression System (ALiCE®):

- 1. In a first purification step, the protein is purified from the cleared cell lysate using StrepTag capture material. Eluate fractions are analyzed by SDS-PAGE.
- Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

Purity:

>80 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.

Endotoxin Level:

Low Endotoxin less than 1 EU/mg (< 0.1 ng/mg)

Grade:

Crystallography grade

Target Details

Target:

DHX16

Alternative Name:

DHX16 (DHX16 Products)

Background:

Pre-mRNA-splicing factor ATP-dependent RNA helicase DHX16 (EC 3.6.4.13) (ATP-dependent RNA helicase #3) (DEAH-box protein 16),FUNCTION: Required for pre-mRNA splicing as component of the spliceosome (PubMed:20423332, PubMed:20841358, PubMed:25296192, PubMed:29360106). Contributes to pre-mRNA splicing after spliceosome formation and prior to the first transesterification reaction. As a component of the minor spliceosome, involved in the splicing of U12-type introns in pre-mRNAs (Probable). Plays also a role in innate antiviral response by acting as a pattern recognition receptor sensing splicing signals in viral RNA (PubMed:35263596). Mechanistically, TRIM6 promotes the interaction between unanchored 'Lys-48'-polyubiquitin chains and DHX16, leading to DHX16 interaction with RIGI and ssRNA to amplify RIGI-dependent innate antiviral immune responses (PubMed:35263596). (ECO:0000269|PubMed:20423332, ECO:0000269|PubMed:20841358, ECO:0000269|PubMed:25296192, ECO:0000269|PubMed:29360106, ECO:0000269|PubMed:35263596, ECO:0000305|PubMed:33509932}.

Molecular Weight:

119.3 kDa

UniProt:

060231

Application Details

Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.
Comment:	ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications. During lysate production, the cell wall and other cellular components that are not required for
	protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer. If you have a special request, please contact us.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	Unlimited (if stored properly)




Image 1. "Crystallography Grade" protein due to multi-step, protein-specific purification process