

# Datasheet for ABIN3092105 DHX9 Protein (AA 1-1270) (Strep Tag)



Overview

Quantity:	250 µg
Target:	DHX9
Protein Characteristics:	AA 1-1270
Origin:	Human
Source:	Cell-free protein synthesis (CFPS)
Protein Type:	Recombinant
Purification tag / Conjugate:	This DHX9 protein is labelled with Strep Tag.
Application:	SDS-PAGE (SDS), ELISA, Western Blotting (WB)

## Product Details

Brand:	AliCE®
Sequence:	MGDVKNFLYA WCGKRKMTPS YEIRAVGNKN RQKFMCEVQV EGYNYTGMGN STNKKDAQSN
	AARDFVNYLV RINEIKSEEV PAFGVASPPP LTDTPDTTAN AEGDLPTTMG GPLPPHLALK
	AENNSEVGAS GYGVPGPTWD RGANLKDYYS RKEEQEVQAT LESEEVDLNA GLHGNWTLEN
	AKARLNQYFQ KEKIQGEYKY TQVGPDHNRS FIAEMTIYIK QLGRRIFARE HGSNKKLAAQ
	SCALSLVRQL YHLGVVEAYS GLTKKKEGET VEPYKVNLSQ DLEHQLQNII QELNLEILPP
	PEDPSVPVAL NIGKLAQFEP SQRQNQVGVV PWSPPQSNWN PWTSSNIDEG PLAFATPEQI
	SMDLKNELMY QLEQDHDLQA ILQERELLPV KKFESEILEA ISQNSVVIIR GATGCGKTTQ
	VPQFILDDFI QNDRAAECNI VVTQPRRISA VSVAERVAFE RGEEPGKSCG YSVRFESILP
	RPHASIMFCT VGVLLRKLEA GIRGISHVIV DEIHERDINT DFLLVVLRDV VQAYPEVRIV
	LMSATIDTSM FCEYFFNCPI IEVYGRTYPV QEYFLEDCIQ MTHFVPPPKD KKKKDKDDDG
	GEDDDANCNL ICGDEYGPET RLSMSQLNEK ETPFELIEAL LKYIETLNVP GAVLVFLPGW

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/6 | Product datasheet for ABIN3092105 | 02/25/2025 | Copyright antibodies-online. All rights reserved. NLIYTMQKHL EMNPHFGSHR YQILPLHSQI PREEQRKVFD PVPVGVTKVI LSTNIAETSI TINDVVYVID SCKQKVKLFT AHNNMTNYAT VWASKTNLEQ RKGRAGRVRP GFCFHLCSRA RFERLETHMT PEMFRTPLHE IALSIKLLRL GGIGQFLAKA IEPPPLDAVI EAEHTLRELD ALDANDELTP LGRILAKLPI EPRFGKMMIM GCIFYVGDAI CTIAAATCFP EPFINEGKRL GYIHRNFAGN RFSDHVALLS VFQAWDDARM GGEEAEIRFC EHKRLNMATL RMTWEAKVQL KEILINSGFP EDCLLTQVFT NTGPDNNLDV VISLLAFGVY PNVCYHKEKR KILTTEGRNA LIHKSSVNCP FSSQDMKYPS PFFVFGEKIR TRAISAKGMT LVTPLQLLLF ASKKVQSDGQ IVLVDDWIKL QISHEAAACI TGLRAAMEAL VVEVTKQPAI ISQLDPVNER MLNMIRQISR PSAAGINLMI GSTRYGDGPR PPKMARYDNG SGYRRGGSSY SGGGYGGGYS SGGYGSGGYG GSANSFRAGY GAGVGGGYRG VSRGGFRGNS GGDYRGPSGG YRGSGGFQRG GGRGAYGTGY FGQGRGGGY

Sequence without tag. The proposed Strep-Tag is based on experience s with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.

#### Characteristics:

Key Benefits:

- Made in Germany from design to production by highly experienced protein experts.
- Protein expressed with ALiCE® and purified in one-step affinity chromatography
- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

#### Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require posttranslational modifications.
- During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system -

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/6 | Product datasheet for ABIN3092105 | 02/25/2025 | Copyright antibodies-online. All rights reserved. all that's needed is the DNA that codes for the desired protein!

#### Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured against its specific reference buffer.
- We use the Expasy's ProtParam tool to determine the absorption coefficient of each protein.

Purification:	One-step Strep-tag purification of proteins expressed in Almost Living Cell-Free Expression System (AliCE®).
Purity:	> 70-80 % as determined by SDS PAGE, Western Blot and analytical SEC (HPLC).
Grade:	custom-made

### Target Details

Target:	DHX9
Alternative Name:	DHX9 (DHX9 Products)
Background:	ATP-dependent RNA helicase A (EC 3.6.4.13) (DEAH box protein 9) (DExH-box helicase 9)
	(Leukophysin) (LKP) (Nuclear DNA helicase II) (NDH II) (RNA helicase A),FUNCTION:
	Multifunctional ATP-dependent nucleic acid helicase that unwinds DNA and RNA in a 3' to 5'
	direction and that plays important roles in many processes, such as DNA replication,
	transcriptional activation, post-transcriptional RNA regulation, mRNA translation and RNA-
	mediated gene silencing (PubMed:9111062, PubMed:11416126, PubMed:12711669,
	PubMed:15355351, PubMed:16680162, PubMed:17531811, PubMed:20669935,
	PubMed:21561811, PubMed:24049074, PubMed:25062910, PubMed:24990949,
	PubMed:28221134). Requires a 3'-single-stranded tail as entry site for acid nuclei unwinding
	activities as well as the binding and hydrolyzing of any of the four ribo- or deoxyribo-nucleotide
	triphosphates (NTPs) (PubMed:1537828). Unwinds numerous nucleic acid substrates such as
	double-stranded (ds) DNA and RNA, DNA:RNA hybrids, DNA and RNA forks composed of either
	partially complementary DNA duplexes or DNA:RNA hybrids, respectively, and also DNA and
	RNA displacement loops (D- and R-loops), triplex-helical DNA (H-DNA) structure and DNA and
	RNA-based G-quadruplexes (PubMed:20669935, PubMed:21561811, PubMed:24049074). Binds
	dsDNA, single-stranded DNA (ssDNA), dsRNA, ssRNA and poly(A)-containing RNA
	(PubMed:9111062, PubMed:10198287). Binds also to circular dsDNA or dsRNA of either linear
	and/or circular forms and stimulates the relaxation of supercoiled DNAs catalyzed by

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/6 | Product datasheet for ABIN3092105 | 02/25/2025 | Copyright antibodies-online. All rights reserved. topoisomerase TOP2A (PubMed:12711669). Plays a role in DNA replication at origins of replication and cell cycle progression (PubMed:24990949). Plays a role as a transcriptional coactivator acting as a bridging factor between polymerase II holoenzyme and transcription factors or cofactors, such as BRCA1, CREBBP, RELA and SMN1 (PubMed:11149922, PubMed:9323138, PubMed:9662397, PubMed:11038348, PubMed:11416126, PubMed:15355351, PubMed:28221134). Binds to the CDKN2A promoter (PubMed:11038348). Plays several roles in post-transcriptional regulation of gene expression (PubMed:28221134, PubMed:28355180). In cooperation with NUP98, promotes pre-mRNA alternative splicing activities of a subset of genes (PubMed:11402034, PubMed:16680162, PubMed:28221134, PubMed:28355180). As component of a large PER complex, is involved in the negative regulation of 3' transcriptional termination of circadian target genes such as PER1 and NR1D1 and the control of the circadian rhythms (By similarity). Acts also as a nuclear resolvase that is able to bind and neutralize harmful massive secondary double-stranded RNA structures formed by inverted-repeat Alu retrotransposon elements that are inserted and transcribed as parts of genes during the process of gene transposition (PubMed:28355180). Involved in the positive regulation of nuclear export of constitutive transport element (CTE)-containing unspliced mRNA (PubMed:9162007, PubMed:10924507, PubMed:11402034). Component of the coding region determinant (CRD)-mediated complex that promotes cytoplasmic MYC mRNA stability (PubMed:19029303). Plays a role in mRNA translation (PubMed:28355180). Positively regulates translation of selected mRNAs through its binding to post-transcriptional control element (PCE) in the 5'-untranslated region (UTR) (PubMed:16680162). Involved with LARP6 in the translation stimulation of type I collagen mRNAs for CO1A1 and CO1A2 through binding of a specific stemloop structure in their 5'-UTRs (PubMed:22190748). Stimulates LIN28A-dependent mRNA translation probably by facilitating ribonucleoprotein remodeling during the process of translation (PubMed:21247876). Plays also a role as a small interfering (siRNA)-loading factor involved in the RNA-induced silencing complex (RISC) loading complex (RLC) assembly, and hence functions in the RISC-mediated gene silencing process (PubMed:17531811). Binds preferentially to short double-stranded RNA, such as those produced during rotavirus intestinal infection (PubMed:28636595). This interaction may mediate NLRP9 inflammasome activation and trigger inflammatory response, including IL18 release and pyroptosis (PubMed:28636595). Finally, mediates the attachment of heterogeneous nuclear ribonucleoproteins (hnRNPs) to actin filaments in the nucleus (PubMed:11687588). {ECO:0000250|UniProtKB:070133, ECO:0000269|PubMed:10198287, ECO:0000269|PubMed:10924507, ECO:0000269|PubMed:11038348, ECO:0000269|PubMed:11149922, ECO:0000269|PubMed:11402034, ECO:0000269|PubMed:11416126, ECO:0000269|PubMed:11687588, ECO:0000269|PubMed:12711669,

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 4/6 | Product datasheet for ABIN3092105 | 02/25/2025 | Copyright antibodies-online. All rights reserved.

ECO:0000269 PubMed:15355351, ECO:0000269 PubMed:1537828,	
ECO:0000269 PubMed:16680162, ECO:0000269 PubMed:17531811,	
EC0:0000269 PubMed:19029303, EC0:0000269 PubMed:20669935,	
ECO:0000269 PubMed:21247876, ECO:0000269 PubMed:21561811,	
EC0:0000269 PubMed:22190748, EC0:0000269 PubMed:24049074,	
EC0:0000269 PubMed:24990949, EC0:0000269 PubMed:25062910,	
EC0:0000269 PubMed:28221134, EC0:0000269 PubMed:28355180,	
ECO:0000269 PubMed:28636595, ECO:0000269 PubMed:9111062,	
ECO:0000269 PubMed:9162007, ECO:0000269 PubMed:9323138,	
ECO:0000269 PubMed:9662397}., FUNCTION: (Microbial infection) Plays a role in HIV-1	
replication and virion infectivity (PubMed:11096080, PubMed:19229320, PubMed:25149208	3,
PubMed:27107641). Enhances HIV-1 transcription by facilitating the binding of RNA	
polymerase II holoenzyme to the proviral DNA (PubMed:11096080, PubMed:25149208). Bin	ıds
(via DRBM domain 2) to the HIV-1 TAR RNA and stimulates HIV-1 transcription of	
transactivation response element (TAR)-containing mRNAs (PubMed:9892698,	
PubMed:11096080). Involved also in HIV-1 mRNA splicing and transport (PubMed:2514920	8).
Positively regulates HIV-1 gag mRNA translation, through its binding to post-transcriptional	
control element (PCE) in the 5'-untranslated region (UTR) (PubMed:16680162). Binds (via	
DRBM domains) to a HIV-1 double-stranded RNA region of the primer binding site (PBS)-	
segment of the 5'-UTR, and hence stimulates DHX9 incorporation into virions and virion	
infectivity (PubMed:27107641). Also plays a role as a cytosolic viral MyD88-dependent DNA	and
RNA sensors in plasmacytoid dendritic cells (pDCs), and hence induce antiviral innate immu	Jne
responses (PubMed:20696886, PubMed:21957149). Binds (via the OB-fold region) to viral	
single-stranded DNA unmethylated C-phosphate-G (CpG) oligonucleotide (PubMed:2069688	36).
{ECO:0000269 PubMed:11096080, ECO:0000269 PubMed:16680162,	
EC0:0000269 PubMed:19229320, EC0:0000269 PubMed:20696886,	
EC0:0000269 PubMed:21957149, EC0:0000269 PubMed:25149208,	
ECO:0000269 PubMed:27107641, ECO:0000269 PubMed:9892698}.	

Molecular Weight:

UniProt:

Q08211

141.0 kDa

# Application Details

Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies
	as well. As the protein has not been tested for functional studies yet we cannot offer a
	guarantee though.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 5/6 | Product datasheet for ABIN3092105 | 02/25/2025 | Copyright antibodies-online. All rights reserved.

## Application Details

Comment:	ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from
	Nicotiana tabacum c.v This contains all the protein expression machinery needed to produce
	even the most difficult-to-express proteins, including those that require post-translational
	modifications.
	During lysate production, the cell wall and other cellular components that are not required for
	protein production are removed, leaving only the protein production machinery and the
	mitochondria to drive the reaction. During our lysate completion steps, the additional
	components needed for protein production (amino acids, cofactors, etc.) are added to produce
	something that functions like a cell, but without the constraints of a living system - all that's
	needed is the DNA that codes for the desired protein!
Restrictions:	For Research Use only

# Handling

Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer. Standard Storage Buffer: PBS pH 7.4, 10 % Glycerol <b>Might differ depending on protein.</b>
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	12 months