

Datasheet for ABIN3092212  
**DYRK2 Protein (AA 1-601) (His tag)**

## 1 Image

[Go to Product page](#)

## Overview

Quantity:	1 mg
Target:	DYRK2
Protein Characteristics:	AA 1-601
Origin:	Human
Source:	Insect Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This DYRK2 protein is labelled with His tag.
Application:	Western Blotting (WB), ELISA, SDS-PAGE (SDS), Crystallization (Crys)

## Product Details

Sequence:	MLTRKPSAAA PAAYPTGRGG DSAVRQLQAS PGLGAGATRS GVGTGPPSPI ALPPLRASNA AAAHTIGGS KHTMNDHLHV GSHAHGQIQV QQLFEDNSNK RTVLTTPNG LTTVGKTGLP VPPERQLDSI HRRQGSSTSL KSMEGMGKVK ATPMTPEQAM KQYMQKLTA EHHEIFSYPE IYFLGLNAKK RQGMTGGPNN GGYDDDQGSY VQVPHDHVAY RYEVCLKVIGK GSFGQVVKAY DHKVHQHVAL KMRVNEKRFH RQAAEEIRIL EHLRKQDKDN TMNVIHMLEN FTFRNHICMT FELLSMNLVE LIKKNKFQGF SLPLVRKFAH SILQCLDALH KNRIIHCCLK PENILLKQQG RSGIKVIDFG SSCYEHQRVY TYIQRFYRA PEVLGARYG MPIDMWSLGC ILAELLTGYP LLPGEDEGDQ LACMIELLGM PSQKLLDASK RAKNFVSSKG YPRYCTVTTL SDGSVVLNNG RSRRGKLRGP PESREWGNAL KGCDDPLFLD FLKQCLEWDP AVRMTPGQAL RHPWLRRLRP KPPTGEKTSV KRITESTGAI TSISKLPPTS SSASKLRTNL AQMTDANGNI QQRTVLPKLV S <b>Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a special request, please contact us.</b>
-----------	---

## Product Details

---

- Characteristics:
- Made in Germany - from design to production - by highly experienced protein experts.
  - Human DYRK2 Protein (raised in Insect Cells) purified by multi-step, protein-specific process to ensure crystallization grade.
  - State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made to order protein and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).

When you order this made-to-order protein you will only pay upon receipt of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered.

The concentration of our recombinant proteins is measured using the absorbance at 280nm.

The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.

The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

- Purification:
- Two step purification of proteins expressed in baculovirus infected SF9 insect cells:
1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.
  2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

Purity: >95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.

Sterility: 0.22 µm filtered

Endotoxin Level: Protein is endotoxin free.

Grade: Crystallography grade

## Target Details

Target:	DYRK2
Alternative Name:	DYRK2 ( <a href="#">DYRK2 Products</a> )
Background:	<p>Serine/threonine-protein kinase involved in the regulation of the mitotic cell cycle, cell proliferation, apoptosis, organization of the cytoskeleton and neurite outgrowth. Functions in part via its role in ubiquitin-dependent proteasomal protein degradation. Functions downstream of ATM and phosphorylates p53/TP53 at 'Ser-46', and thereby contributes to the induction of apoptosis in response to DNA damage. Phosphorylates NFATC1, and thereby inhibits its accumulation in the nucleus and its transcription factor activity. Phosphorylates EIF2B5 at 'Ser-544', enabling its subsequent phosphorylation and inhibition by GSK3B. Likewise, phosphorylation of NFATC1, CRMP2/DPYSL2 and CRMP4/DPYSL3 promotes their subsequent phosphorylation by GSK3B. May play a general role in the priming of GSK3 substrates. Inactivates GYS1 by phosphorylation at 'Ser-641', and potentially also a second phosphorylation site, thus regulating glycogen synthesis. Mediates EDVP E3 ligase complex formation and is required for the phosphorylation and subsequent degradation of KATNA1. Phosphorylates TERT at 'Ser-457', promoting TERT ubiquitination by the EDVP complex. Phosphorylates SIAH2, and thereby increases its ubiquitin ligase activity. Promotes the proteasomal degradation of MYC and JUN, and thereby regulates progress through the mitotic cell cycle and cell proliferation. Promotes proteasomal degradation of GLI2 and GLI3, and thereby plays a role in smoothened and sonic hedgehog signaling. Plays a role in cytoskeleton organization and neurite outgrowth via its phosphorylation of DCX and DPYSL2. Phosphorylates CRMP2/DPYSL2, CRMP4/DPYSL3, DCX, EIF2B5, EIF4EBP1, GLI2, GLI3, GYS1, JUN, MDM2, MYC, NFATC1, p53/TP53, TAU/MAPT and KATNA1. Can phosphorylate histone H1, histone H3 and histone H2B (in vitro). Can phosphorylate CARHSP1 (in vitro).</p> <p>{ECO:0000269 PubMed:11311121, ECO:0000269 PubMed:12588975, ECO:0000269 PubMed:14593110, ECO:0000269 PubMed:15910284, ECO:0000269 PubMed:16511445, ECO:0000269 PubMed:16611631, ECO:0000269 PubMed:17349958, ECO:0000269 PubMed:18455992, ECO:0000269 PubMed:18599021, ECO:0000269 PubMed:19287380, ECO:0000269 PubMed:22307329, ECO:0000269 PubMed:22878263, ECO:0000269 PubMed:23362280, ECO:0000269 PubMed:9748265}.</p>
Molecular Weight:	67.6 kDa Including tag.
UniProt:	<a href="#">Q92630</a>
Pathways:	<a href="#">Regulation of Carbohydrate Metabolic Process</a>

## Application Details

Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.
Comment:	In cases in which it is highly likely that the recombinant protein with the default tag will be insoluble our protein lab may suggest a higher molecular weight tag (e.g. GST-tag) instead to increase solubility. We will discuss all possible options with you in detail to assure that you receive your protein of interest.
Restrictions:	For Research Use only

## Handling

Format:	Liquid
Buffer:	100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	Unlimited (if stored properly)

## Images



**Image 1.** „Crystallography Grade“ protein due to multi-step, protein-specific purification process