antibodies .- online.com





GCN2 Protein (AA 1-1649) (His tag)



Image



Go to Product page

Overview

Quantity:	1 mg
Target:	GCN2 (EIF2AK4)
Protein Characteristics:	AA 1-1649
Origin:	Human
Source:	Insect Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This GCN2 protein is labelled with His tag.
Application:	Western Blotting (WB), SDS-PAGE (SDS), ELISA, Crystallization (Crys)

Product Details

Sequence:

MAGGRGAPGR GRDEPPESYP QRQDHELQAL EAIYGADFQD LRPDACGPVK EPPEINLVLY
PQGLTGEEVY VKVDLRVKCP PTYPDVVPEI ELKNAKGLSN ESVNLLKSRL EELAKKHCGE
VMIFELAYHV QSFLSEHNKP PPKSFHEEML ERRAQEEQQR LLEAKRKEEQ EQREILHEIQ
RRKEEIKEEK KRKEMAKQER LEIASLSNQD HTSKKDPGGH RTAAILHGGS PDFVGNGKHR
ANSSGRSRRE RQYSVCNSED SPGSCEILYF NMGSPDQLMV HKGKCIGSDE QLGKLVYNAL
ETATGGFVLL YEWVLQWQKK MGPFLTSQEK EKIDKCKKQI QGTETEFNSL VKLSHPNVVR
YLAMNLKEQD DSIVVDILVE HISGVSLAAH LSHSGPIPVH QLRRYTAQLL SGLDYLHSNS
VVHKVLSASN VLVDAEGTVK ITDYSISKRL ADICKEDVFE QTRVRFSDNA LPYKTGKKGD
VWRLGLLLLS LSQGQECGEY PVTIPSDLPA DFQDFLKKCV CLDDKERWSP QQLLKHSFIN
PQPKMPLVEQ SPEDSEGQDY VETVIPSNRL PSAAFFSETQ RQFSRYFIEF EELQLLGKGA
FGAVIKVQNK LDGCCYAVKR IPINPASRQF RRIKGEVTLL SRLHHENIVR YYNAWIERHE
RPAGPGTPPP DSGPLAKDDR AARGQPASDT DGLDSVEAAA PPPILSSSVE WSTSGERSAS

ARFPATGPGS SDDEDDDEDE HGGVFSQSFL PASDSESDII FDNEDENSKS QNQDEDCNEK NGCHESEPSV TTEAVHYLYI QMEYCEKSTL RDTIDQGLYR DTVRLWRLFR EILDGLAYIH EKGMIHRDLK PVNIFLDSDD HVKIGDFGLA TDHLAFSADS KQDDQTGDLI KSDPSGHLTG MVGTALYVSP EVQGSTKSAY NQKVDLFSLG IIFFEMSYHP MVTASERIFV LNQLRDPTSP KFPEDFDDGE HAKQKSVISW LLNHDPAKRP TATELLKSEL LPPPQMEESE LHEVLHHTLT NVDGKAYRTM MAQIFSQRIS PAIDYTYDSD ILKGNFSIRT AKMQQHVCET IIRIFKRHGA VOLCTPLLLP RNROIYEHNE AALFMDHSGM LVMLPFDLRI PFARYVARNN ILNLKRYCIE RVFRPRKLDR FHPKELLECA FDIVTSTTNS FLPTAEIIYT IYEIIQEFPA LQERNYSIYL NHTMLLKAIL LHCGIPEDKL SQVYIILYDA VTEKLTRREV EAKFCNLSLS SNSLCRLYKF IEQKGDLQDL MPTINSLIKQ KTGIAQLVKY GLKDLEEVVG LLKKLGIKLQ VLINLGLVYK VQQHNGIIFQ FVAFIKRRQR AVPEILAAGG RYDLLIPQFR GPQALGPVPT AIGVSIAIDK ISAAVLNMEE SVTISSCDLL VVSVGQMSMS RAINLTQKLW TAGITAEIMY DWSQSQEELQ EYCRHHEITY VALVSDKEGS HVKVKSFEKE RQTEKRVLET ELVDHVLQKL RTKVTDERNG REASDNLAVQ NLKGSFSNAS GLFEIHGATV VPIVSVLAPE KLSASTRRRY ETQVQTRLQT SLANLHQKSS EIEILAVDLP KETILQFLSL EWDADEQAFN TTVKQLLSRL PKQRYLKLVC DEIYNIKVEK KVSVLFLYSY RDDYYRIL

Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a special request, please contact us.

Characteristics:

- Made in Germany from design to production by highly experienced protein experts.
- Human EIF2AK4 Protein (raised in Insect Cells) purified by multi-step, protein-specific process to ensure crystallization grade.
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made to order protein and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).

When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered.

The concentration of our recombinant proteins is measured using the absorbance at 280nm.

The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.

The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

Purification:

Two step purification of proteins expressed in baculovirus infected SF9 insect cells:

- 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.
- 2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

Purity:

>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.

Sterility:

0.22 µm filtered

Endotoxin Level:

Protein is endotoxin free.

Grade:

Crystallography grade

Target Details

Target:

GCN2 (EIF2AK4)

Alternative Name:

EIF2AK4 (EIF2AK4 Products)

Background:

Metabolic-stress sensing protein kinase that phosphorylates the alpha subunit of eukaryotic translation initiation factor 2 (eIF-2-alpha/EIF2S1) on 'Ser-52' in response to low amino acid availability (PubMed:25329545). Plays a role as an activator of the integrated stress response (ISR) required for adaptation to amino acid starvation. Converts phosphorylated eIF-2-alpha/EIF2S1 either to a competitive inhibitor of the translation initiation factor eIF-2B, leading to a global protein synthesis repression, and thus to a reduced overall utilization of amino acids, or to a translational initiation activation of specific mRNAs, such as the transcriptional activator ATF4, and hence allowing ATF4-mediated reprogramming of amino acid biosynthetic gene expression to alleviate nutrient depletion. Binds uncharged tRNAs (By similarity). Involved in cell cycle arrest by promoting cyclin D1 mRNA translation repression after the unfolded protein response pathway (UPR) activation or cell cycle inhibitor CDKN1A/p21 mRNA translation activation in response to amino acid deprivation (PubMed:26102367). Plays a role in the consolidation of synaptic plasticity, learning as well as formation of long-term memory. Plays a role in neurite outgrowth inhibition. Plays a proapoptotic role in response to glucose

Expiry Date:

	deprivation. Promotes global cellular protein synthesis repression in response to UV irradiation
	independently of the stress-activated protein kinase/c-Jun N-terminal kinase (SAPK/JNK) and
	p38 MAPK signaling pathways (By similarity). Plays a role in the antiviral response against
	alphavirus infection, impairs early viral mRNA translation of the incoming genomic virus RNA,
	thus preventing alphavirus replication (By similarity). {ECO:0000250 UniProtKB:P15442,
	ECO:0000250 UniProtKB:Q9QZ05, ECO:0000269 PubMed:25329545,
	ECO:0000269 PubMed:26102367}., (Microbial infection) Plays a role in modulating the adaptive
	immune response to yellow fever virus infection, promotes dendritic cells to initiate autophagy
	and antigene presentation to both CD4(+) and CD8(+) T-cells under amino acid starvation
	(PubMed:24310610). {ECO:0000269 PubMed:24310610}.
Molecular Weight:	187.7 kDa Including tag.
UniProt:	Q9P2K8
Pathways:	ER-Nucleus Signaling, Hepatitis C
Application Details	
Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies
	as well. As the protein has not been tested for functional studies yet we cannot offer a gurantee
	though.
Comment:	In cases in which it is highly likely that the recombinant protein with the default tag will be
	insoluble our protein lab may suggest a higher molecular weight tag (e.g. GST-tag) instead to
	increase solubility. We will discuss all possible options with you in detail to assure that you
	receive your protein of interest.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.

Unlimited (if stored properly)



Image 1. "Crystallography Grade" protein due to multi-step, protein-specific purification process