antibodies .- online.com





Ephrin A5 Protein (EFNA5) (AA 21-203) (His tag)



()	1/0	r\ /1	014	
()	ve	I V I	-v	V

Overview	
Quantity:	1 mg
Target:	Ephrin A5 (EFNA5)
Protein Characteristics:	AA 21-203
Origin:	Human
Source:	Insect Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This Ephrin A5 protein is labelled with His tag.
Application:	SDS-PAGE (SDS), Western Blotting (WB), ELISA, Crystallization (Crys)
Product Details	
Sequence:	QDPGSKAVAD RYAVYWNSSN PRFQRGDYHI DVCINDYLDV FCPHYEDSVP EDKTERYVLY
	MVNFDGYSAC DHTSKGFKRW ECNRPHSPNG PLKFSEKFQL FTPFSLGFEF RPGREYFYIS
	SAIPDNGRRS CLKLKVFVRP TNSCMKTIGV HDRVFDVNDK VENSLEPADD TVHESAEPSR GEN
	Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a
	special request, please contact us.
Characteristics:	 Made in Germany - from design to production - by highly experienced protein experts. Human EFNA5 Protein (raised in Insect Cells) purified by multi-step, protein-specific process to ensure crystallization grade. State-of-the-art algorithm used for plasmid design (Gene synthesis).
	This protein is a made to order protein and will be made for the first time for your order. Our
	experts in the lab will ensure that you receive a correctly folded protein.
	The big advantage of ordering our made-to-order proteins in comparison to ordering custom

made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).

When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered.

The concentration of our recombinant proteins is measured using the absorbance at 280nm.

The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.

The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

Purification:

Two step purification of proteins expressed in baculovirus infected SF9 insect cells:

- 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.
- 2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

Purity:

>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.

Sterility:

0.22 µm filtered

Endotoxin Level:

Protein is endotoxin free.

Grade:

Crystallography grade

Target Details

Target:	Ephrin A5 (EFNA5)	
Alternative Name:	EFNA5 (EFNA5 Products)	
Background:	Cell surface GPI-bound ligand for Eph receptors, a family of receptor tyrosine kinases which a	
	crucial for migration, repulsion and adhesion during neuronal, vascular and epithelial	
	development. Binds promiscuously Eph receptors residing on adjacent cells, leading to contact-	
	dependent bidirectional signaling into neighboring cells. The signaling pathway downstream of	

the receptor is referred to as forward signaling while the signaling pathway downstream of the
ephrin ligand is referred to as reverse signaling. Induces compartmentalized signaling within a
caveolae-like membrane microdomain when bound to the extracellular domain of its cognate
receptor. This signaling event requires the activity of the Fyn tyrosine kinase. Activates the
EPHA3 receptor to regulate cell-cell adhesion and cytoskeletal organization. With the receptor
EPHA2 may regulate lens fiber cells shape and interactions and be important for lens
transparency maintenance. May function actively to stimulate axon fasciculation. The
interaction of EFNA5 with EPHA5 also mediates communication between pancreatic islet cells
to regulate glucose-stimulated insulin secretion. Cognate/functional ligand for EPHA7, their
interaction regulates brain development modulating cell-cell adhesion and repulsion.
{ECO:0000269 PubMed:10601038, ECO:0000269 PubMed:11870224}.

Molecular Weight:	22.1 kDa Including tag.	
UniProt:	P52803	
Pathways:	RTK Signaling	

Application Details

Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies	
	as well. As the protein has not been tested for functional studies yet we cannot offer a gurantee	
	though.	
Comment:	In cases in which it is highly likely that the recombinant protein with the default tag will be	
	insoluble our protein lab may suggest a higher molecular weight tag (e.g. GST-tag) instead to	
	increase solubility. We will discuss all possible options with you in detail to assure that you	
	receive your protein of interest.	
Restrictions:	For Research Use only	

Handling

Format:	Liquid	
Buffer:	100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer.	
Handling Advice:	Avoid repeated freeze-thaw cycles.	
Storage:	-80 °C	
Storage Comment:	Store at -80°C.	

1	1 11	١٠
\vdash	land	unc
	ıarıa	11110

Expiry Date:

Unlimited (if stored properly)