

Datasheet for ABIN3092274 EIF3L Protein (AA 2-564) (His tag)



Go to Product page

Overview

Quantity:	1 mg	
Target:	EIF3L	
Protein Characteristics:	AA 2-564	
Origin:	Human	
Source:	Escherichia coli (E. coli)	
Protein Type:	Recombinant	
Purification tag / Conjugate:	This EIF3L protein is labelled with His tag.	
Application:	ELISA, Western Blotting (WB), Crystallization (Crys), SDS-PAGE (SDS)	

Product Details

Sequence:

SYPADDYESE AAYDPYAYPS DYDMHTGDPK QDLAYERQYE QQTYQVIPEV IKNFIQYFHK
TVSDLIDQKV YELQASRVSS DVIDQKVYEI QDIYENSWTK LTERFFKNTP WPEAEAIAPQ
VGNDAVFLIL YKELYYRHIY AKVSGGPSLE QRFESYYNYC NLFNYILNAD GPAPLELPNQ
WLWDIIDEFI YQFQSFSQYR CKTAKKSEEE IDFLRSNPKI WNVHSVLNVL HSLVDKSNIN
RQLEVYTSGG DPESVAGEYG RHSLYKMLGY FSLVGLLRLH SLLGDYYQAI KVLENIELNK
KSMYSRVPEC QVTTYYYVGF AYLMMRRYQD AIRVFANILL YIQRTKSMFQ RTTYKYEMIN
KQNEQMHALL AIALTMYPMR IDESIHLQLR EKYGDKMLRM QKGDPQVYEE LFSYSCPKFL
SPVVPNYDNV HPNYHKEPFL QQLKVFSDEV QQQAQLSTIR SFLKLYTTMP VAKLAGFLDL
TEQEFRIQLL VFKHKMKNLV WTSGISALDG EFQSASEVDF YIDKDMIHIA DTKVARRYGD
FFIRQIHKFE ELNRTLKKMG QRP

Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a special request, please contact us.

Characteristics:

- Made in Germany from design to production by highly experienced protein experts.
- Human EIF3L Protein (raised in E. Coli) purified by multi-step, protein-specific process to ensure crystallization grade.
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made to order protein and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).

When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered.

The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.

The concentration of our recombinant proteins is measured using the absorbance at 280nm.

The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

Purification:

Two step purification of proteins expressed in bacterial culture:

- 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.
- 2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

Purity:	>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.
Sterility:	0.22 μm filtered
Endotoxin Level:	Endotoxin has not been removed. Please contact us if you require endotoxin removal.
Grade:	Crystallography grade

Target Details

Target:	EIF3L		
Alternative Name:	EIF3L (EIF3L Products)		
Background:	Component of the eukaryotic translation initiation factor 3 (eIF-3) complex, which is required for several steps in the initiation of protein synthesis. The eIF-3 complex associates with the 40S ribosome and facilitates the recruitment of eIF-1, eIF-1A, eIF-2:GTP:methionyl-tRNAi and eIF-5 to form the 43S pre-initiation complex (43S PIC). The eIF-3 complex stimulates mRNA recruitment to the 43S PIC and scanning of the mRNA for AUG recognition. The eIF-3 complex		
	is also required for disassembly and recycling of post-termination ribosomal complexes and subsequently prevents premature joining of the 40S and 60S ribosomal subunits prior to initiation. In case of FCV infection, plays a role in the ribosomal termination-reinitiation event leading to the translation of VP2 (PubMed:18056426). {ECO:0000269 PubMed:18056426}.		
Molecular Weight:	67.5 kDa Including tag.		
UniProt:	Q9Y262		
Pathways:	Ribonucleoprotein Complex Subunit Organization		
Application Details			
Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a gurantee though.		
Comment:	In cases in which it is highly likely that the recombinant protein with the default tag will be insoluble our protein lab may suggest a higher molecular weight tag (e.g. GST-tag) instead to increase solubility. We will discuss all possible options with you in detail to assure that you receive your protein of interest.		
Restrictions:	For Research Use only		
Handling			
Format:	Liquid		
Buffer:	100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer.		
Handling Advice:	Avoid repeated freeze-thaw cycles.		
Storage:	-80 °C		
Storage Comment:	Store at -80°C.		

1	1 11	١٠
\vdash	land	∥n∩
1	ıarıa	11110

Expiry Date:

Unlimited (if stored properly)