

Datasheet for ABIN3092656

GABBR1 Protein (AA 855-961) (His tag)



Overviev	

Overview		
Quantity:	1 mg	
Target:	GABBR1	
Protein Characteristics:	AA 855-961	
Origin:	Human	
Source:	Insect Cells	
Protein Type:	Recombinant	
Purification tag / Conjugate:	This GABBR1 protein is labelled with His tag.	
Application:	Western Blotting (WB), SDS-PAGE (SDS), ELISA, Crystallization (Crys)	
Product Details		
Sequence:	PKMRRLITRG EWQSEAQDTM KTGSSTNNNE EEKSRLLEKE NRELEKIIAE KEERVSELRH	
	QLQSRQQLRS RRHPPTPPEP SGGLPRGPPE PPDRLSCDGS RVHLLYK	
	Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a	
	special request, please contact us.	
Characteristics:	 Made in Germany - from design to production - by highly experienced protein experts. Human GABBR1 Protein (raised in Insect Cells) purified by multi-step, protein-specific process to ensure crystallization grade. State-of-the-art algorithm used for plasmid design (Gene synthesis). 	
	This protein is a made to order protein and will be made for the first time for your order. Our	
	experts in the lab will ensure that you receive a correctly folded protein.	
	The big advantage of ordering our made-to-order proteins in comparison to ordering custom	
	made proteins from other companies is that there is no financial obligation in case the protein	

cannot be expressed or purified.

In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).

When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered. The concentration of our recombinant proteins is measured using the absorbance at 280nm. The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.

The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

Purification:

Two step purification of proteins expressed in baculovirus infected SF9 insect cells:

- 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.
- 2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

Purity:

>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.

Sterility:

0.22 µm filtered

GABBR1

Endotoxin Level:

Protein is endotoxin free.

Grade:

Target:

Crystallography grade

Target Details

Alternative Name:	GABBR1 (GABBR1 Products)
Background:	Component of a heterodimeric G-protein coupled receptor for GABA, formed by GABBR1 and
	GABBR2. Within the heterodimeric GABA receptor, only GABBR1 seems to bind agonists, while
	GABBR2 mediates coupling to G proteins. Ligand binding causes a conformation change that
	triggers signaling via guanine nucleotide-binding proteins (G proteins) and modulates the
	activity of down-stream effectors, such as adenylate cyclase. Signaling inhibits adenylate

cyclase, stimulates phospholipase A2, activates potassium channels, inactivates voltage-dependent calcium-channels and modulates inositol phospholipid hydrolysis. Calcium is required for high affinity binding to GABA. Plays a critical role in the fine-tuning of inhibitory synaptic transmission. Pre-synaptic GABA receptor inhibits neurotransmitter release by down-regulating high-voltage activated calcium channels, whereas postsynaptic GABA receptor decreases neuronal excitability by activating a prominent inwardly rectifying potassium (Kir) conductance that underlies the late inhibitory postsynaptic potentials. Not only implicated in synaptic inhibition but also in hippocampal long-term potentiation, slow wave sleep, muscle relaxation and antinociception. Activated by (-)-baclofen, cgp27492 and blocked by phaclofen., Isoform 1E may regulate the formation of functional GABBR1/GABBR2 heterodimers by competing for GABBR2 binding. This could explain the observation that certain small molecule ligands exhibit differential affinity for central versus peripheral sites.

Molecular Weight:	13.4 kDa Including tag.	
UniProt:	Q9UBS5	
Pathways:	Positive Regulation of Peptide Hormone Secretion, cAMP Metabolic Process	

Application Details

Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies	
	as well. As the protein has not been tested for functional studies yet we cannot offer a gurantee	
	though.	

In cases in which it is highly likely that the recombinant protein with the default tag will be insoluble our protein lab may suggest a higher molecular weight tag (e.g. GST-tag) instead to increase solubility. We will discuss all possible options with you in detail to assure that you receive your protein of interest.

Restrictions: For Research Use only

Handling

Comment:

Format:	Liquid	
Buffer:	100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer.	
Handling Advice:	Avoid repeated freeze-thaw cycles.	
Storage:	-80 °C	
Storage Comment:	Store at -80°C.	

\vdash	land	linc
	iaria	11116

Expiry Date:

Unlimited (if stored properly)