

Datasheet for ABIN3092786

GNAI1 Protein (AA 2-354) (His tag)[Go to Product page](#)**1** Image

Overview

| | |
|-------------------------------|--|
| Quantity: | 1 mg |
| Target: | GNAI1 |
| Protein Characteristics: | AA 2-354 |
| Origin: | Human |
| Source: | Insect Cells |
| Protein Type: | Recombinant |
| Purification tag / Conjugate: | This GNAI1 protein is labelled with His tag. |
| Application: | ELISA, Western Blotting (WB), SDS-PAGE (SDS), Crystallization (Crys) |

Product Details

| | |
|------------------|--|
| Sequence: | <p>GCTLSAEDKA AVERSKMIDR NLREDGEKAA REVKLLLLGA GESGKSTIVK QMKIIHEAGY SEEECKQYKA VVYSNTIQSI IAIIRAMGRL KIDFGDSARA DDARQLFVLA GAAEEGFMTA ELAGVIKRLW KDSGVQACFN RSREYQLNDS AAYYLNDLDR IAQPNYIPTQ QDVLRTRVKT TGIVETHFTF KDLHFKMFDV GGQRSERKKW IHCFEVGTAI IFCVALSDYD LVLAEDEEMN RMHESMKLFD SICNNKWFTD TSILFLNKK DLFEEDIKKS PLTICYPEYA GSNTYEEAAA YIQCQFEDLN KRKDTKEIYT HFTCATDTKN VQFVFDAVTD VIKNNLKDC GLF</p> <p>Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a special request, please contact us.</p> |
| Characteristics: | <ul style="list-style-type: none">• Made in Germany - from design to production - by highly experienced protein experts.• Human GNAI1 Protein (raised in Insect Cells) purified by multi-step, protein-specific process to ensure crystallization grade.• State-of-the-art algorithm used for plasmid design (Gene synthesis). |

Product Details

This protein is a made to order protein and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).

When you order this made-to-order protein you will only pay upon receipt of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered.

The concentration of our recombinant proteins is measured using the absorbance at 280nm. The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.

The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

| | |
|------------------|--|
| Purification: | Two step purification of proteins expressed in baculovirus infected SF9 insect cells: <ol style="list-style-type: none">1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot. |
| Purity: | >95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot. |
| Sterility: | 0.22 µm filtered |
| Endotoxin Level: | Protein is endotoxin free. |
| Grade: | Crystallography grade |

Target Details

| | |
|-------------------|--|
| Target: | GNAI1 |
| Alternative Name: | GNAI1 (GNAI1 Products) |
| Background: | Guanine nucleotide-binding proteins (G proteins) function as transducers downstream of G |

Target Details

protein-coupled receptors (GPCRs) in numerous signaling cascades. The alpha chain contains the guanine nucleotide binding site and alternates between an active, GTP-bound state and an inactive, GDP-bound state. Signaling by an activated GPCR promotes GDP release and GTP binding. The alpha subunit has a low GTPase activity that converts bound GTP to GDP, thereby terminating the signal. Both GDP release and GTP hydrolysis are modulated by numerous regulatory proteins (PubMed:8774883, PubMed:18434541). Signaling is mediated via effector proteins, such as adenylate cyclase. Inhibits adenylate cyclase activity, leading to decreased intracellular cAMP levels (By similarity). The inactive GDP-bound form prevents the association of RGS14 with centrosomes and is required for the translocation of RGS14 from the cytoplasm to the plasma membrane. Required for normal cytokinesis during mitosis (PubMed:17635935). {ECO:0000250|UniProtKB:P10824, ECO:0000269|PubMed:17635935, ECO:0000269|PubMed:18434541, ECO:0000269|PubMed:8774883}.

Molecular Weight: 41.2 kDa Including tag.

UniProt: [P63096](#)

Pathways: [G-protein mediated Events](#)

Application Details

Application Notes: In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.

Comment: In cases in which it is highly likely that the recombinant protein with the default tag will be insoluble our protein lab may suggest a higher molecular weight tag (e.g. GST-tag) instead to increase solubility. We will discuss all possible options with you in detail to assure that you receive your protein of interest.

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: 100 mM NaCl, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer.

Handling Advice: Avoid repeated freeze-thaw cycles.

Storage: -80 °C

Storage Comment: Store at -80°C.

Handling

Expiry Date: Unlimited (if stored properly)

Images



Image 1. „Crystallography Grade“ protein due to multi-step, protein-specific purification process