
antibodies -online.com





GRIA2 Protein (AA 25-543) (His tag)



Go to Product page

Overview

Quantity:	1 mg	
Target:	GRIA2	
Protein Characteristics:	AA 25-543	
Origin:	Human	
Source:	Insect Cells	
Protein Type:	Recombinant	
Purification tag / Conjugate:	This GRIA2 protein is labelled with His tag.	
Application:	Western Blotting (WB), SDS-PAGE (SDS), Crystallization (Crys), ELISA	

Product Details

Sequence:

NSIQIGGLFP RGADQEYSAF RVGMVQFSTS EFRLTPHIDN LEVANSFAVT NAFCSQFSRG VYAIFGFYDK KSVNTITSFC GTLHVSFITP SFPTDGTHPF VIQMRPDLKG ALLSLIEYYQ WDKFAYLYDS DRGLSTLQAV LDSAAEKKWQ VTAINVGNIN NDKKDEMYRS LFQDLELKKE RRVILDCERD KVNDIVDQVI TIGKHVKGYH YIIANLGFTD GDLLKIQFGG ANVSGFQIVD YDDSLVSKFI ERWSTLEEKE YPGAHTTTIK YTSALTYDAV QVMTEAFRNL RKQRIEISRR GNAGDCLANP AVPWGQGVEI ERALKQVQVE GLSGNIKFDQ NGKRINYTIN IMELKTNGPR KIGYWSEVDK MVVTLTELPS GNDTSGLENK TVVVTTILES PYVMMKKNHE MLEGNERYEG YCVDLAAEIA KHCGFKYKLT IVGDGKYGAR DADTKIWNGM VGELVYGKAD IAIAPLTITL VREEVIDFSK PFMSLGISIM IKKPQKSKPG VFSFLDPLA

Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a special request, please contact us.

Characteristics:

- Made in Germany from design to production by highly experienced protein experts.
- Human GRIA2 Protein (raised in Insect Cells) purified by multi-step, protein-specific process to ensure crystallization grade.
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made to order protein and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).

When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered.

The concentration of our recombinant proteins is measured using the absorbance at 280nm.

The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.

The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

Purification:

Two step purification of proteins expressed in baculovirus infected SF9 insect cells:

- 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.
- 2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

Purity: >95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.

Sterility: 0.22 µm filtered

Endotoxin Level: Protein is endotoxin free.

Grade: Crystallography grade

Target Details

-		
Target:	GRIA2	
Alternative Name:	GRIA2 (GRIA2 Products)	
Background:	Receptor for glutamate that functions as ligand-gated ion channel in the central nervous system and plays an important role in excitatory synaptic transmission. L-glutamate acts as an excitatory neurotransmitter at many synapses in the central nervous system. Binding of the excitatory neurotransmitter L-glutamate induces a conformation change, leading to the opening of the cation channel, and thereby converts the chemical signal to an electrical impulse. The receptor then desensitizes rapidly and enters a transient inactive state, characterized by the presence of bound agonist. In the presence of CACNG4 or CACNG7 or CACNG8, shows resensitization which is characterized by a delayed accumulation of current flux upon continued	
Molecular Weight:	application of glutamate. {ECO:0000269 PubMed:20614889}. 59.2 kDa Including tag.	
UniProt:	P42262	
Pathways:	PI3K-Akt Signaling	
Application Details		
Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a gurantee though.	
Comment:	In cases in which it is highly likely that the recombinant protein with the default tag will be insoluble our protein lab may suggest a higher molecular weight tag (e.g. GST-tag) instead to increase solubility. We will discuss all possible options with you in detail to assure that you receive your protein of interest.	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Buffer:	100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer.	
Handling Advice:	Avoid repeated freeze-thaw cycles.	
Storage:	-80 °C	
Storage Comment:	Store at -80°C.	

\vdash	land	linc
	iaria	11116

Expiry Date:

Unlimited (if stored properly)