antibodies .- online.com





HAP1 Protein (AA 1-671) (His tag)



Image



Go to Product page

Overview

Quantity:	1 mg
Target:	HAP1
Protein Characteristics:	AA 1-671
Origin:	Human
Source:	Insect Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This HAP1 protein is labelled with His tag.
Application:	SDS-PAGE (SDS), Western Blotting (WB), ELISA, Crystallization (Crys)

Product Details

Sequence:

MRPKRLGRCC AGSRLGPGDP AALTCAPSPS ASPAPEPSAQ PQARGTGQRV GSRATSGSQF LSEARTGARP ASEAGAKAGA RRPSAFSAIQ GDVRSMPDNS DAPWTRFVFQ GPFGSRATGR GTGKAAGIWK TPAAYVGRRP GVSGPERAAF IRELEEALCP NLPPPVKKIT QEDVKVMLYL LEELLPPVWE SVTYGMVLQR ERDLNTAARI GQSLVKQNSV LMEENSKLEA LLGSAKEEIL YLRHQVNLRD ELLQLYSDSD EEDEDEEEEE EEKEAEEEQE EEEAEEDLQC AHPCDAPKLI SQEALLHQHH CPQLEALQEK LRLLEEENHQ LREEASQLDT LEDEEQMLIL ECVEQFSEAS QQMAELSEVL VLRLENYERQ QQEVARLQAQ VLKLQQRCRM YGAETEKLQK QLASEKEIQM QLQEESVWVG SQLQDLREKY MDCGGMLIEM QEEVKTLRQQ PPVSTGSATH YPYSVPLETL PGFQETLAEE LRTSLRRMIS DPVYFMERNY EMPRGDTSSL RYDFRYSEDR EQVRGFEAEE GLMLAADIMR GEDFTPAEEF VPQEELGAAK KVPAEEGVME EAELVSEETE GWEEVELELD EATRMNVVTS ALEASGLGPS HLDMNYVLQQ LANWQDAHYR RQLRWKMLQK GECPHGALPA ASRTSCRSSC R

Endotoxin Level:

Grade:

Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a special request, please contact us. Characteristics: Made in Germany - from design to production - by highly experienced protein experts. Human HAP1 Protein (raised in Insect Cells) purified by multi-step, protein-specific process to ensure crystallization grade. • State-of-the-art algorithm used for plasmid design (Gene synthesis). This protein is a made to order protein and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein. The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified. In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization). When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered. The concentration of our recombinant proteins is measured using the absorbance at 280nm. The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer. The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein. Purification: Two step purification of proteins expressed in baculovirus infected SF9 insect cells: 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE. 2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot. >95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot. Purity: Sterility: 0.22 µm filtered

Protein is endotoxin free.

Crystallography grade

Target Details

Target:	HAP1
Alternative Name:	HAP1 (HAP1 Products)
Background:	Originally identified as neuronal protein that specifically associates with HTT/huntingtin and the
	binding is enhanced by an expanded polyglutamine repeat within HTT possibly affecting HAP1
	interaction properties. Both HTT and HAP1 are involved in intracellular trafficking and HAP1 is
	proposed to link HTT to motor proteins and/or transport cargos. Seems to play a role in
	vesicular transport within neurons and axons such as from early endosomes to late endocytic
	compartments and to promote neurite outgrowth. The vesicular transport function via
	association with microtubule-dependent transporters can be attenuated by association with
	mutant HTT. Involved in the axonal transport of BDNF and its activity-dependent secretion, the
	function seems to involve HTT, DCTN1 and a complex with SORT1. Involved in APP trafficking
	and seems to faciltate APP anterograde transport and membrane insertion thereby possibly
	reducing processing into amyloid beta. Involved in delivery of gamma-aminobutyric acid
	(GABA(A)) receptors to synapses, the function is dependent on kinesin motor protein KIF5 and
	is disrupted by HTT with expanded polyglutamine repeat. Involved in regulation of
	autophagosome motility by promoting efficient retrograde axonal transport. Seems to be
	involved in regulation of membrane receptor recycling and degradation, and respective signal
	transduction, including GABA(A) receptors, tyrosine kinase receptors, EGFR, IP3 receptor and
	androgen receptor. Among others suggested to be involved in control of feeding behavior
	(involving hypothalamic GABA(A) receptors), cerebellar and brainstem development (involving
	AHI1 and NTRK1/TrkA), postnatal neurogenesis (involving hypothalamic NTRK2/TrkB), and
	ITPR1/InsP3R1-mediated Ca(2+) release (involving HTT and possibly the effect of mutant
	HTT). Via association with DCTN1/dynactin p150-glued and HTT/huntingtin involved in
	cytoplasmic retention of REST in neurons. May be involved in ciliogenesis. Involved in
	regulation of exocytosis. Seems to be involved in formation of cytoplasmic inclusion bodies
	(STBs). In case of anomalous expression of TBP, can sequester a subset of TBP into STBs,
	sequestration is enhanced by an expanded polyglutamine repeat within TBP. HAP1-containing
	STBs have been proposed to play a protective role against neurodegeneration in Huntigton
	disease (HD) and spinocerebellar ataxia 17 (SCA17). {ECO:0000269 PubMed:18922795}.
Molecular Weight:	76.5 kDa Including tag.
UniProt:	P54257
Pathways:	Cell RedoxHomeostasis, Smooth Muscle Cell Migration, Positive Regulation of Response to

DNA Damage Stimulus

Application Details

Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a gurantee though.
Comment:	In cases in which it is highly likely that the recombinant protein with the default tag will be insoluble our protein lab may suggest a higher molecular weight tag (e.g. GST-tag) instead to increase solubility. We will discuss all possible options with you in detail to assure that you receive your protein of interest.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	Unlimited (if stored properly)

Images




Image 1. "Crystallography Grade" protein due to multi-step, protein-specific purification process