

Datasheet for ABIN3093023

IKZF1 Protein (AA 1-519) (Strep Tag)



[Go to Product page](#)

Overview

Quantity:	250 µg
Target:	IKZF1
Protein Characteristics:	AA 1-519
Origin:	Human
Source:	Cell-free protein synthesis (CFPS)
Protein Type:	Recombinant
Purification tag / Conjugate:	This IKZF1 protein is labelled with Strep Tag.
Application:	ELISA, Western Blotting (WB), SDS-PAGE (SDS)

Product Details

Brand:	AliCE®
Sequence:	<p>MDADEGQDMS QVSGKESPPV SDTPDEGDEP MPIPEDLSTT SGGQQSSKSD RVVASNVKVE TQSDEENGRA CEMNGEECAE DLRMLDASGE KMNGSHRDQG SSALSGVGGI RLPNGKLKCD ICGIICIGPN VLMVHKRSHT GERPFQCNQC GASFTQKGNL LRHIKLHSGE KPFKCHLCNY ACRRRDALTG HLRTHSVGKP HKCGYCGRSY QQRSSLEEHK ERCHNYLESM GLPGTLYPVI KEETNHSEMA EDLCKIGSER SLVLDRLASN VAKRKSSMPQ KFLGDKGLSD TPYDSSASYE KENEMMKSHV MDQAINNAIN YLGAESLRPL VQTPPGGSEV VPVISPMYQL HKPLAEGTPR SNHSAQDSAV ENLLLLSKAK LVPSEREASP SNSCQDSTDT ESNNEEQRSQ LIYLTNHIAP HARNGLSLKE EHRAYDLLRA ASENSQDALR VVSTSGEQMK VYKCEHCRVL FLDHVMYTIH MGCHGFRDPF ECNMCGYHSQ DRYEFSSHIT RGEHRFHMS</p> <p>Sequence without tag. The proposed Strep-Tag is based on experience s with the expression system, a different complexity of the protein could make another tag necessary. In case you</p>

have a special request, please contact us.

Characteristics:

Key Benefits:

- Made in Germany - from design to production - by highly experienced protein experts.
- Protein expressed with ALiCE® and purified in one-step affinity chromatography
- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from *Nicotiana tabacum* c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.
- During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured against its specific reference buffer.
- We use the ExPASy's ProtParam tool to determine the absorption coefficient of each protein.

Purification:

One-step Strep-tag purification of proteins expressed in Almost Living Cell-Free Expression System (ALiCE®).

Purity:

> 70-80 % as determined by SDS PAGE, Western Blot and analytical SEC (HPLC).

Grade:

custom-made

Target Details

Target:	IKZF1
Alternative Name:	IKZF1 (IKZF1 Products)
Background:	<p>DNA-binding protein Ikaros (Ikaros family zinc finger protein 1) (Lymphoid transcription factor LyF-1),FUNCTION: Transcription regulator of hematopoietic cell differentiation (PubMed:17934067). Binds gamma-satellite DNA (PubMed:17135265, PubMed:19141594). Plays a role in the development of lymphocytes, B- and T-cells. Binds and activates the enhancer (delta-A element) of the CD3-delta gene. Repressor of the TDT (fikzfterminal deoxynucleotidyltransferase) gene during thymocyte differentiation. Regulates transcription through association with both HDAC-dependent and HDAC-independent complexes. Targets the 2 chromatin-remodeling complexes, NuRD and BAF (SWI/SNF), in a single complex (PYR complex), to the beta-globin locus in adult erythrocytes. Increases normal apoptosis in adult erythroid cells. Confers early temporal competence to retinal progenitor cells (RPCs) (By similarity). Function is isoform-specific and is modulated by dominant-negative inactive isoforms (PubMed:17135265, PubMed:17934067). {ECO:0000250 UniProtKB:Q03267, ECO:0000269 PubMed:10204490, ECO:0000269 PubMed:17135265, ECO:0000269 PubMed:17934067, ECO:0000269 PubMed:19141594}.</p>
Molecular Weight:	57.5 kDa
UniProt:	Q13422
Pathways:	Production of Molecular Mediator of Immune Response

Application Details

Application Notes:	<p>In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.</p>
Comment:	<p>ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from <i>Nicotiana tabacum</i> c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.</p> <p>During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's</p>

Application Details

	needed is the DNA that codes for the desired protein!
Restrictions:	For Research Use only

Handling

Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer. Standard Storage Buffer: PBS pH 7.4, 10 % Glycerol Might differ depending on protein.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	12 months