antibodies -online.com





CLNS1A Protein (AA 2-237) (His tag)



| iew |
|-----|
| |

| Quantity: | 1 mg |
|-------------------------------|--|
| Target: | CLNS1A |
| Protein Characteristics: | AA 2-237 |
| Origin: | Human |
| Source: | Escherichia coli (E. coli) |
| Protein Type: | Recombinant |
| Purification tag / Conjugate: | This CLNS1A protein is labelled with His tag. |
| Application: | ELISA, Western Blotting (WB), SDS-PAGE (SDS), Crystallization (Crys) |
| Product Details | |
| Sequence: | SFLKSFPPPG PAEGLLRQQP DTEAVLNGKG LGTGTLYIAE SRLSWLDGSG LGFSLEYPTI |
| | SLHALSRDRS DCLGEHLYVM VNAKFEEESK EPVADEEEED SDDDVEPITE FRFVPSDKSA |
| | LEAMFTAMCE CQALHPDPED EDSDDYDGEE YDVEAHEQGQ GDIPTFYTYE EGLSHLTAEG |
| | QATLERLEGM LSQSVSSQYN MAGVRTEDSI RDYEDGMEVD TTPTVAGQFE DADVDH |
| | Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a |
| | special request, please contact us. |
| Characteristics: | Made in Germany - from design to production - by highly experienced protein experts. Human CLNS1A Protein (raised in E. Coli) purified by multi-step, protein-specific process to ensure crystallization grade. State-of-the-art algorithm used for plasmid design (Gene synthesis). |
| | This protein is a made to order protein and will be made for the first time for your order. Our |
| | experts in the lab will ensure that you receive a correctly folded protein. |

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).

When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered. The concentration of our recombinant proteins is measured using the absorbance at 280nm.

The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.

The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

Purification:

Two step purification of proteins expressed in bacterial culture:

- In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.
- 2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

Purity:

>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.

Sterility:

0.22 µm filtered

Endotoxin Level:

Endotoxin has not been removed. Please contact us if you require endotoxin removal.

Grade:

Crystallography grade

Target Details

| Target: | CLNS1A |
|-------------------|--|
| Alternative Name: | CLNS1A (CLNS1A Products) |
| Background: | Chaperone that regulates the assembly of spliceosomal U1, U2, U4 and U5 small nuclear ribonucleoproteins (snRNPs), the building blocks of the spliceosome. Thereby, plays an |
| | important role in the splicing of cellular pre-mRNAs. Most spliceosomal snRNPs contain a |

| ECO:0000269 PubMed:18984161}. |
|---|
| {ECO:0000269 PubMed:10330151, ECO:0000269 PubMed:11713266, |
| activation of a swelling-induced chloride conductance pathway. |
| and their transport to the nucleus. May also indirectly participate in cellular volume control by |
| Sm proteins and their transfer to an SMN-Sm complex triggers the assembly of core snRNPs |
| assembly of the core snRNP. Dissociation by the SMN complex of CLNS1A from the trapped |
| trapped in an inactive 6S pICIn-Sm complex by the chaperone CLNS1A that controls the |
| core snRNP. In the cytosol, the Sm proteins SNRPD1, SNRPD2, SNRPE, SNRPF and SNRPG are |
| that assemble in a heptameric protein ring on the Sm site of the small nuclear RNA to form the |
| common set of Sm proteins SNRPB, SNRPD1, SNRPD2, SNRPD3, SNRPE, SNRPF and SNRPG |

| Molecular Weight: | 27.0 kDa Including tag. |
|-------------------|--|
| UniProt: | P54105 |
| Pathways: | Ribonucleoprotein Complex Subunit Organization |

Application Details

Application Notes:

| | as well. As the protein has not been tested for functional studies yet we cannot offer a gurantee though. |
|---------------|---|
| Comment: | In cases in which it is highly likely that the recombinant protein with the default tag will be insoluble our protein lab may suggest a higher molecular weight tag (e.g. GST-tag) instead to increase solubility. We will discuss all possible options with you in detail to assure that you receive your protein of interest. |
| Restrictions: | For Research Use only |

In addition to the applications listed above we expect the protein to work for functional studies

Handling

| Format: | Liquid |
|------------------|--|
| Buffer: | 100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer. |
| Handling Advice: | Avoid repeated freeze-thaw cycles. |
| Storage: | -80 °C |
| Storage Comment: | Store at -80°C. |
| Expiry Date: | Unlimited (if stored properly) |