
antibodies .- online.com





CAMK2D Protein (AA 2-499) (His tag)



Image



Go to Product page

Overview

Quantity:	1 mg
Target:	CAMK2D
Protein Characteristics:	AA 2-499
Origin:	Human
Source:	Insect Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This CAMK2D protein is labelled with His tag.
Application:	ELISA, Western Blotting (WB), SDS-PAGE (SDS), Crystallization (Crys)

Product Details

Sequence:

ASTTTCTRFT DEYQLFEELG KGAFSVVRRC MKIPTGQEYA AKIINTKKLS ARDHQKLERE
ARICRLLKHP NIVRLHDSIS EEGFHYLVFD LVTGGELFED IVAREYYSEA DASHCIQQIL
ESVNHCHLNG IVHRDLKPEN LLLASKSKGA AVKLADFGLA IEVQGDQQAW FGFAGTPGYL
SPEVLRKDPY GKPVDMWACG VILYILLVGY PPFWDEDQHR LYQQIKAGAY DFPSPEWDTV
TPEAKDLINK MLTINPAKRI TASEALKHPW ICQRSTVASM MHRQETVDCL KKFNARRKLK
GAILTTMLAT RNFSAAKSLL KKPDGVKEST ESSNTTIEDE DVKARKQEII KVTEQLIEAI
NNGDFEAYTK ICDPGLTAFE PEALGNLVEG MDFHRFYFEN ALSKSNKPIH TIILNPHVHL
VGDDAACIAY IRLTQYMDGS GMPKTMQSEE TRVWHRRDGK WQNVHFHRSG SPTVPIKPPC
IPNGKENFSG GTSLWQNI

Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a special request, please contact us.

Characteristics:

- Made in Germany from design to production by highly experienced protein experts.
- Human CAMK2D Protein (raised in Insect Cells) purified by multi-step, protein-specific process to ensure crystallization grade.
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made to order protein and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).

When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered.

The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.

The concentration of our recombinant proteins is measured using the absorbance at 280nm.

The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

Purification:

Two step purification of proteins expressed in baculovirus infected SF9 insect cells:

- 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.
- 2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

Purity:

>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.

Sterility:

0.22 µm filtered

Endotoxin Level:

Protein is endotoxin free.

Grade:

Crystallography grade

Target Details

Target:	CAMK2D
Alternative Name:	CAMK2D (CAMK2D Products)
Background:	Calcium/calmodulin-dependent protein kinase involved in the regulation of Ca(2+) homeostatis
	and excitation-contraction coupling (ECC) in heart by targeting ion channels, transporters and
	accessory proteins involved in Ca(2+) influx into the myocyte, Ca(2+) release from the
	sarcoplasmic reticulum (SR), SR Ca(2+) uptake and Na(+) and K(+) channel transport. Targets
	also transcription factors and signaling molecules to regulate heart function. In its activated
	form, is involved in the pathogenesis of dilated cardiomyopathy and heart failure. Contributes to
	cardiac decompensation and heart failure by regulating SR Ca(2+) release via direct
	phosphorylation of RYR2 Ca(2+) channel on 'Ser-2808'. In the nucleus, phosphorylates the
	MEF2 repressor HDAC4, promoting its nuclear export and binding to 14-3-3 protein, and
	expression of MEF2 and genes involved in the hypertrophic program. Is essential for left
	ventricular remodeling responses to myocardial infarction. In pathological myocardial
	remodeling acts downstream of the beta adrenergic receptor signaling cascade to regulate key
	proteins involved in ECC. Regulates Ca(2+) influx to myocytes by binding and phosphorylating
	the L-type Ca(2+) channel subunit beta-2 CACNB2. In addition to Ca(2+) channels, can target
	and regulate the cardiac sarcolemmal Na(+) channel Nav1.5/SCN5A and the K+ channel
	Kv4.3/KCND3, which contribute to arrhythmogenesis in heart failure. Phosphorylates
	phospholamban (PLN/PLB), an endogenous inhibitor of SERCA2A/ATP2A2, contributing to the
	enhancement of SR Ca(2+) uptake that may be important in frequency-dependent acceleration
	of relaxation (FDAR) and maintenance of contractile function during acidosis. May participate in
	the modulation of skeletal muscle function in response to exercise, by regulating SR Ca(2+)
	transport through phosphorylation of PLN/PLB and triadin, a ryanodine receptor-coupling
	factor. {ECO:0000269 PubMed:16690701, ECO:0000269 PubMed:17179159}.
Molecular Weight:	57.2 kDa Including tag.
UniProt:	Q13557
Pathways:	WNT Signaling, Interferon-gamma Pathway, Myometrial Relaxation and Contraction, Smooth
	Muscle Cell Migration
Application Details	
Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies
	as well. As the protein has not been tested for functional studies yet we cannot offer a gurantee
	though.

Application Details

Images

Comment:	In cases in which it is highly likely that the recombinant protein with the default tag will be insoluble our protein lab may suggest a higher molecular weight tag (e.g. GST-tag) instead to increase solubility. We will discuss all possible options with you in detail to assure that you receive your protein of interest.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	Unlimited (if stored properly)




Image 1. "Crystallography Grade" protein due to multi-step, protein-specific purification process