

Datasheet for ABIN3093234 IST1 Protein (AA 1-364) (Strep Tag)



Go to Product page

_					
	W	0	rv	10	W

Quantity:	250 μg
Target:	IST1
Protein Characteristics:	AA 1-364
Origin:	Human
Source:	Cell-free protein synthesis (CFPS)
Protein Type:	Recombinant
Purification tag / Conjugate:	This IST1 protein is labelled with Strep Tag.
Application:	ELISA, SDS-PAGE (SDS), Western Blotting (WB)

Product Details

Brand:	AliCE®
Sequence:	MLGSGFKAER LRVNLRLVIN RLKLLEKKKT ELAQKARKEI ADYLAAGKDE RARIRVEHII
	REDYLVEAME ILELYCDLLL ARFGLIQSMK ELDSGLAESV STLIWAAPRL QSEVAELKIV
	ADQLCAKYSK EYGKLCRTNQ IGTVNDRLMH KLSVEAPPKI LVERYLIEIA KNYNVPYEPD
	SVVMAEAPPG VETDLIDVGF TDDVKKGGPG RGGSGGFTAP VGGPDGTVPM PMPMPMPSAN
	TPFSYPLPKG PSDFNGLPMG TYQAFPNIHP PQIPATPPSY ESVDDINADK NISSAQIVGP
	GPKPEASAKL PSRPADNYDN FVLPELPSVP DTLPTASAGA STSASEDIDF DDLSRRFEEL KKKT
	Sequence without tag. The proposed Strep-Tag is based on experience s with the expression
	system, a different complexity of the protein could make another tag necessary. In case you
	have a special request, please contact us.
Characteristics:	Key Benefits:

- Made in Germany from design to production by highly experienced protein experts.
- · Protein expressed with ALiCE® and purified in one-step affinity chromatography
- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- · State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require posttranslational modifications.
- During lysate production, the cell wall and other cellular components that are not required for
 protein production are removed, leaving only the protein production machinery and the
 mitochondria to drive the reaction. During our lysate completion steps, the additional
 components needed for protein production (amino acids, cofactors, etc.) are added to
 produce something that functions like a cell, but without the constraints of a living system all that's needed is the DNA that codes for the desired protein!

Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured against its specific reference buffer.
- · We use the Expasy's ProtParam tool to determine the absorption coefficient of each protein.

Purification:	One-step Strep-tag purification of proteins expressed in Almost Living Cell-Free Expression System (AliCE®).
Purity:	> 70-80 % as determined by SDS PAGE, Western Blot and analytical SEC (HPLC).
Grade:	custom-made
Target Details	
Target:	IST1

Target Details

Alternative Name:	IST1 (IST1 Products)
Background:	IST1 homolog (hIST1) (Charged multivesicular body protein 8) (CHMP8) (Putative MAPK-
	activating protein PM28),FUNCTION: ESCRT-III-like protein involved in cytokinesis, nuclear
	envelope reassembly and endosomal tubulation (PubMed:19129479, PubMed:26040712,
	PubMed:28242692). Is required for efficient abscission during cytokinesis (PubMed:19129479)
	Involved in recruiting VPS4A and/or VPS4B to the midbody of dividing cells (PubMed:19129480
	PubMed:19129479). During late anaphase, involved in nuclear envelope reassembly and mitotic
	spindle disassembly together with the ESCRT-III complex: IST1 acts by mediating the
	recruitment of SPAST to the nuclear membrane, leading to microtubule severing
	(PubMed:26040712). Recruited to the reforming nuclear envelope (NE) during anaphase by
	LEMD2 (PubMed:28242692). Regulates early endosomal tubulation together with the ESCRT-III
	complex by mediating the recruitment of SPAST (PubMed:23897888).
	{ECO:0000269 PubMed:19129479, ECO:0000269 PubMed:19129480,
	ECO:0000269 PubMed:23897888, ECO:0000269 PubMed:26040712,
	ECO:0000269 PubMed:28242692}.
Molecular Weight:	39.8 kDa
UniProt:	P53990
Pathways:	Regulation of Cell Size
Application Details	
Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies
	as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.
Comment:	ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from
	Nicotiana tabacum c.v This contains all the protein expression machinery needed to produce
	even the most difficult-to-express proteins, including those that require post-translational modifications.
	During lysate production, the cell wall and other cellular components that are not required for
	protein production are removed, leaving only the protein production machinery and the
	mitochondria to drive the reaction. During our lysate completion steps, the additional

Application Details

Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer. Standard Storage Buffer: PBS pH 7.4, 10 % Glycerol Might differ depending on protein.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	12 months