antibodies -online.com





KISS1 Protein (AA 20-138) (His tag)



Image



Go to Product page

Overview

1
KI
aracteristics: A
H
In
pe: Re
n tag / Conjugate: Th
n: W
Details
EF
SF
Se
sp
stics: •
Tł
ех
Tł
m
Hi In Dee: Re Thag / Conjugate: Th W Details EF SF Se Se Th

cannot be expressed or purified.

In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).

When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered. The concentration of our recombinant proteins is measured using the absorbance at 280nm. The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.

The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

Purification:

Two step purification of proteins expressed in baculovirus infected SF9 insect cells:

- 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.
- 2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

Purity:

>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.

Sterility:

0.22 µm filtered

Endotoxin Level:

Protein is endotoxin free.

Grade:

Crystallography grade

Target Details

Target:	KISS1
Alternative Name:	KISS1 (KISS1 Products)
Background:	Metastasis suppressor protein in malignant melanomas and in some breast cancers. May
	regulate events downstream of cell-matrix adhesion, perhaps involving cytoskeletal
	reorganization. Generates a C-terminally amidated peptide, metastin which functions as the
	endogenous ligand of the G-protein coupled receptor GPR54. Activation of the receptor inhibits
	cell proliferation and cell migration, key characteristics of tumor metastasis. Kp-10 is a

Expiry Date:	Unlimited (if stored properly)
Storage Comment:	Store at -80°C.
Storage:	-80 °C
Handling Advice:	Avoid repeated freeze-thaw cycles.
Buffer:	100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer.
Format:	Liquid
Handling	
Restrictions:	For Research Use only
Comment:	In cases in which it is highly likely that the recombinant protein with the default tag will be insoluble our protein lab may suggest a higher molecular weight tag (e.g. GST-tag) instead to increase solubility. We will discuss all possible options with you in detail to assure that you receive your protein of interest.
Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a gurantee though.
Application Details	
Pathways:	Positive Regulation of Peptide Hormone Secretion
UniProt:	Q15726
Molecular Weight:	13.5 kDa Including tag.
	in isolated first trimester trophoblasts. Kp-10 is a paracrine/endocrine regulator in fine-tuning trophoblast invasion generated by the trophoblast itself. The receptor is also essential for normal gonadotropin-released hormone physiology and for puberty. The hypothalamic KiSS1/GPR54 system is a pivotal factor in central regulation of the gonadotropic axis at puberty and in adulthood. {ECO:0000269 PubMed:11060311, ECO:0000269 PubMed:11385580, ECO:0000269 PubMed:15500545, ECO:0000269 PubMed:9185708}.
	decapeptide derived from the primary translation product, isolated in conditioned medium of first trimester trophoblast. Kp-10, but not other kisspeptins, increased intracellular Ca(2+) levels



Image 1. "Crystallography Grade" protein due to multi-step, protein-specific purification process