antibodies .- online.com





PKC delta Protein (AA 1-329) (His tag)



Overview

Quantity:	1 mg
Target:	PKC delta (PKCd)
Protein Characteristics:	AA 1-329
Origin:	Human
Source:	Insect Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This PKC delta protein is labelled with His tag.
Application:	Western Blotting (WB), SDS-PAGE (SDS), ELISA, Crystallization (Crys)

Product Details

Sequence:

MAPFLRIAFN SYELGSLQAE DEANQPFCAV KMKEALSTER GKTLVQKKPT MYPEWKSTFD
AHIYEGRVIQ IVLMRAAEEP VSEVTVGVSV LAERCKKNNG KAEFWLDLQP QAKVLMSVQY
FLEDVDCKQS MRSEDEAKFP TMNRRGAIKQ AKIHYIKNHE FIATFFGQPT FCSVCKDFVW
GLNKQGYKCR QCNAAIHKKC IDKIIGRCTG TAANSRDTIF QKERFNIDMP HRFKVHNYMS
PTFCDHCGSL LWGLVKQGLK CEDCGMNVHH KCREKVANLC GINQKLLAEA LNQVTQRASR
RSDSASSEPV GIYQGFEKKT GVAGEDMQD

Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a special request, please contact us.

Characteristics:

- Made in Germany from design to production by highly experienced protein experts.
- Human PRKCD Protein (raised in Insect Cells) purified by multi-step, protein-specific process to ensure crystallization grade.
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made to order protein and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).

When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered.

The concentration of our recombinant proteins is measured using the absorbance at 280nm.

The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.

The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

Purification:

Two step purification of proteins expressed in baculovirus infected SF9 insect cells:

- 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.
- Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

Purity:

>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.

Sterility:

0.22 µm filtered

Endotoxin Level:

Protein is endotoxin free.

Grade:

Crystallography grade

Target Details

Target:	PKC delta (PKCd)
Alternative Name:	PRKCD (PKCd Products)
Background:	Calcium-independent, phospholipid- and diacylglycerol (DAG)-dependent serine/threonine-

protein kinase that plays contrasting roles in cell death and cell survival by functioning as a proapoptotic protein during DNA damage-induced apoptosis, but acting as an anti-apoptotic protein during cytokine receptor-initiated cell death, is involved in tumor suppression as well as survival of several cancers, is required for oxygen radical production by NADPH oxidase and acts as positive or negative regulator in platelet functional responses. Negatively regulates B cell proliferation and also has an important function in self-antigen induced B cell tolerance induction. Upon DNA damage, activates the promoter of the death-promoting transcription factor BCLAF1/Btf to trigger BCLAF1-mediated p53/TP53 gene transcription and apoptosis. In response to oxidative stress, interact with and activate CHUK/IKKA in the nucleus, causing the phosphorylation of p53/TP53. In the case of ER stress or DNA damage-induced apoptosis, can form a complex with the tyrosine-protein kinase ABL1 which trigger apoptosis independently of p53/TP53. In cytosol can trigger apoptosis by activating MAPK11 or MAPK14, inhibiting AKT1 and decreasing the level of X-linked inhibitor of apoptosis protein (XIAP), whereas in nucleus induces apoptosis via the activation of MAPK8 or MAPK9. Upon ionizing radiation treatment, is required for the activation of the apoptosis regulators BAX and BAK, which trigger the mitochondrial cell death pathway. Can phosphorylate MCL1 and target it for degradation which is sufficient to trigger for BAX activation and apoptosis. Is required for the control of cell cycle progression both at G1/S and G2/M phases. Mediates phorbol 12-myristate 13-acetate (PMA)induced inhibition of cell cycle progression at G1/S phase by up-regulating the CDK inhibitor CDKN1A/p21 and inhibiting the cyclin CCNA2 promoter activity. In response to UV irradiation can phosphorylate CDK1, which is important for the G2/M DNA damage checkpoint activation. Can protect glioma cells from the apoptosis induced by TNFSF10/TRAIL, probably by inducing increased phosphorylation and subsequent activation of AKT1. Is highly expressed in a number of cancer cells and promotes cell survival and resistance against chemotherapeutic drugs by inducing cyclin D1 (CCND1) and hyperphosphorylation of RB1, and via several pro-survival pathways, including NF-kappa-B, AKT1 and MAPK1/3 (ERK1/2). Can also act as tumor suppressor upon mitogenic stimulation with PMA or TPA. In N-formyl-methionyl-leucylphenylalanine (fMLP)-treated cells, is required for NCF1 (p47-phox) phosphorylation and activation of NADPH oxidase activity, and regulates TNF-elicited superoxide anion production in neutrophils, by direct phosphorylation and activation of NCF1 or indirectly through MAPK1/3 (ERK1/2) signaling pathways. May also play a role in the regulation of NADPH oxidase activity in eosinophil after stimulation with IL5, leukotriene B4 or PMA. In collagen-induced platelet aggregation, acts a negative regulator of filopodia formation and actin polymerization by interacting with and negatively regulating VASP phosphorylation. Downstream of PAR1, PAR4 and CD36/GP4 receptors, regulates differentially platelet dense granule secretion, acts as a positive regulator in PAR-mediated granule secretion, whereas it negatively regulates

	CD36/GP4-mediated granule release. Phosphorylates MUC1 in the C-terminal and regulates the interaction between MUC1 and beta-catenin. The catalytic subunit phosphorylates 14-3-3 proteins (YWHAB, YWHAZ and YWHAH) in a sphingosine-dependent fashion (By similarity).	
	Phosphorylates ELAVL1 in response to angiotensin-2 treatment (PubMed:18285462).	
	{ECO:0000250, ECO:0000269 PubMed:11748588, ECO:0000269 PubMed:11877440,	
	ECO:0000269 PubMed:15774464, ECO:0000269 PubMed:16940418,	
	ECO:0000269 PubMed:18285462, ECO:0000269 PubMed:19587372,	
	ECO:0000269 PubMed:19801500}.	
Molecular Weight:	38.3 kDa Including tag.	
UniProt:	Q05655	
Pathways:	Interferon-gamma Pathway, EGFR Signaling Pathway, Neurotrophin Signaling Pathway, Thyroid	
	Hormone Synthesis, Regulation of Actin Filament Polymerization, Carbohydrate Homeostasis,	
	Myometrial Relaxation and Contraction, M Phase, G-protein mediated Events, Dicarboxylic Acid	
	Transport, Positive Regulation of Response to DNA Damage Stimulus, Interaction of EGFR with	
	phospholipase C-gamma, Thromboxane A2 Receptor Signaling, Lipid Metabolism	
Application Details		
Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies	
	as well. As the protein has not been tested for functional studies yet we cannot offer a gurantee	
	though.	
Comment:	In cases in which it is highly likely that the recombinant protein with the default tag will be	
	insoluble our protein lab may suggest a higher molecular weight tag (e.g. GST-tag) instead to	
	increase solubility. We will discuss all possible options with you in detail to assure that you	
	receive your protein of interest.	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Buffer:	100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer.	
Handling Advice:	Avoid repeated freeze-thaw cycles.	
Storage:	-80 °C	
Storage Comment:	Store at -80°C.	

\vdash	land	linc
	iaria	11116

Expiry Date:

Unlimited (if stored properly)