

## Datasheet for ABIN3093598

# LIM Domain Kinase 1 Protein (LIMK1) (AA 1-647) (Strep Tag)



Go to Product page

## Overview

Quantity:	250 μg
Target:	LIM Domain Kinase 1 (LIMK1)
Protein Characteristics:	AA 1-647
Origin:	Human
Source:	Cell-free protein synthesis (CFPS)
Protein Type:	Recombinant
Purification tag / Conjugate:	This LIM Domain Kinase 1 protein is labelled with Strep Tag.
Application:	ELISA, Western Blotting (WB), SDS-PAGE (SDS)

Brand:	AliCE®
Sequence:	MRLTLLCCTW REERMGEEGS ELPVCASCGQ RIYDGQYLQA LNADWHADCF RCCDCSASLS
	HQYYEKDGQL FCKKDYWARY GESCHGCSEQ ITKGLVMVAG ELKYHPECFI CLTCGTFIGD
	GDTYTLVEHS KLYCGHCYYQ TVVTPVIEQI LPDSPGSHLP HTVTLVSIPA SSHGKRGLSV
	SIDPPHGPPG CGTEHSHTVR VQGVDPGCMS PDVKNSIHVG DRILEINGTP IRNVPLDEID
	LLIQETSRLL QLTLEHDPHD TLGHGLGPET SPLSSPAYTP SGEAGSSARQ KPVLRSCSID
	RSPGAGSLGS PASQRKDLGR SESLRVVCRP HRIFRPSDLI HGEVLGKGCF GQAIKVTHRE
	TGEVMVMKEL IRFDEETQRT FLKEVKVMRC LEHPNVLKFI GVLYKDKRLN FITEYIKGGT
	LRGIIKSMDS QYPWSQRVSF AKDIASGMAY LHSMNIIHRD LNSHNCLVRE NKNVVVADFG
	LARLMVDEKT QPEGLRSLKK PDRKKRYTVV GNPYWMAPEM INGRSYDEKV DVFSFGIVLC
	EIIGRVNADP DYLPRTMDFG LNVRGFLDRY CPPNCPPSFF PITVRCCDLD PEKRPSFVKL
	EHWLETLRMH LAGHLPLGPQ LEQLDRGFWE TYRRGESGLP AHPEVPD

Sequence without tag. The proposed Strep-Tag is based on experience s with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.

#### Characteristics:

#### Key Benefits:

- · Made in Germany from design to production by highly experienced protein experts.
- · Protein expressed with ALiCE® and purified in one-step affinity chromatography
- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

#### Expression System:

- ALICE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require posttranslational modifications.
- During lysate production, the cell wall and other cellular components that are not required for
  protein production are removed, leaving only the protein production machinery and the
  mitochondria to drive the reaction. During our lysate completion steps, the additional
  components needed for protein production (amino acids, cofactors, etc.) are added to
  produce something that functions like a cell, but without the constraints of a living system all that's needed is the DNA that codes for the desired protein!

#### Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured against its specific reference buffer.
- · We use the Expasy's ProtParam tool to determine the absorption coefficient of each protein.

Purification:	One-step Strep-tag purification of proteins expressed in Almost Living Cell-Free Expression System (AliCE®).
Purity:	> 70-80 % as determined by SDS PAGE, Western Blot and analytical SEC (HPLC).

## **Product Details** Grade: custom-made Target Details Target: LIM Domain Kinase 1 (LIMK1) Alternative Name LIMK1 (LIMK1 Products) Background: LIM domain kinase 1 (LIMK-1) (EC 2.7.11.1), FUNCTION: Serine/threonine-protein kinase that plays an essential role in the regulation of actin filament dynamics. Acts downstream of several Rho family GTPase signal transduction pathways (PubMed:10436159, PubMed:11832213, PubMed:12807904, PubMed:15660133, PubMed:16230460, PubMed:18028908, PubMed:22328514, PubMed:23633677). Activated by upstream kinases including ROCK1, PAK1 and PAK4, which phosphorylate LIMK1 on a threonine residue located in its activation loop (PubMed:10436159). LIMK1 subsequently phosphorylates and inactivates the actin binding/depolymerizing factors cofilin-1/CFL1, cofilin-2/CFL2 and destrin/DSTN, thereby preventing the cleavage of filamentous actin (F-actin), and stabilizing the actin cytoskeleton (PubMed:11832213, PubMed:15660133, PubMed:16230460, PubMed:23633677). In this way LIMK1 regulates several actin-dependent biological processes including cell motility, cell cycle progression, and differentiation (PubMed:11832213, PubMed:15660133, PubMed:16230460, PubMed:23633677). Phosphorylates TPPP on serine residues, thereby promoting microtubule disassembly (PubMed:18028908). Stimulates axonal outgrowth and may be involved in brain development (PubMed:18028908). {ECO:0000269|PubMed:10436159, ECO:0000269|PubMed:11832213, ECO:0000269|PubMed:12807904, ECO:0000269|PubMed:15660133, ECO:0000269|PubMed:16230460, ECO:0000269|PubMed:18028908, ECO:0000269|PubMed:22328514, ECO:0000269|PubMed:23633677}., FUNCTION: [Isoform 3]: Has a dominant negative effect on actin cytoskeletal changes. Required for atypical chemokine receptor ACKR2-induced phosphorylation of cofilin (CFL1). {ECO:0000269|PubMed:10196227}. Molecular Weight: 72.6 kDa UniProt: P53667

Caspase Cascade in Apoptosis, Regulation of Cell Size, CXCR4-mediated Signaling Events

## **Application Details**

Pathways:

In addition to the applications listed above we expect the protein to work for functional studies **Application Notes:** as well. As the protein has not been tested for functional studies yet we cannot offer a

### **Application Details**

Expiry Date:

Application Details	
	guarantee though.
Comment:	ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from
	Nicotiana tabacum c.v This contains all the protein expression machinery needed to produce
	even the most difficult-to-express proteins, including those that require post-translational
	modifications.
	During lysate production, the cell wall and other cellular components that are not required for
	protein production are removed, leaving only the protein production machinery and the
	mitochondria to drive the reaction. During our lysate completion steps, the additional
	components needed for protein production (amino acids, cofactors, etc.) are added to produce
	something that functions like a cell, but without the constraints of a living system - all that's
	needed is the DNA that codes for the desired protein!
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer.
	Standard Storage Buffer: PBS pH 7.4, 10 % Glycerol Might differ depending on protein.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.

12 months