# antibodies .- online.com





### MKNK2 Protein (AA 1-465) (Strep Tag)

2 Images



Go to Product page

#### Overview

Quantity:	1 mg
Target:	MKNK2
Protein Characteristics:	AA 1-465
Origin:	Human
Source:	Tobacco (Nicotiana tabacum)
Protein Type:	Recombinant
Purification tag / Conjugate:	This MKNK2 protein is labelled with Strep Tag.
Application:	ELISA, Western Blotting (WB), SDS-PAGE (SDS)

#### **Product Details**

Sequence:

MVQKKPAELQ GFHRSFKGQN PFELAFSLDQ PDHGDSDFGL QCSARPDMPA SQPIDIPDAK KRGKKKKRGR ATDSFSGRFE DVYQLQEDVL GEGAHARVQT CINLITSQEY AVKIIEKQPG HIRSRVFREV EMLYQCQGHR NVLELIEFFE EEDRFYLVFE KMRGGSILSH IHKRRHFNEL EASVVVQDVA SALDFLHNKG IAHRDLKPEN ILCEHPNQVS PVKICDFDLG SGIKLNGDCS PISTPELLTP CGSAEYMAPE VVEAFSEEAS IYDKRCDLWS LGVILYILLS GYPPFVGRCG SDCGWDRGEA CPACQNMLFE SIQEGKYEFP DKDWAHISCA AKDLISKLLV RDAKQRLSAA QVLQHPWVQG CAPENTLPTP MVLQRNSCAK DLTSFAAEAI AMNRQLAQHD EDLAEEEAAG QGQPVLVRAT SRCLQLSPPS QSKLAQRRQR ASLSSAPVVL VGDHA

Sequence without tag. The proposed Strep-Tag is based on experience s with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.

Characteristics:

Key Benefits:

- · Made in Germany from design to production by highly experienced protein experts.
- Protein expressed with ALiCE® and purified by multi-step, protein-specific process to ensure correct folding and modification.
- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

#### Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require posttranslational modifications.
- During lysate production, the cell wall and other cellular components that are not required for
  protein production are removed, leaving only the protein production machinery and the
  mitochondria to drive the reaction. During our lysate completion steps, the additional
  components needed for protein production (amino acids, cofactors, etc.) are added to
  produce something that functions like a cell, but without the constraints of a living system all that's needed is the DNA that codes for the desired protein!

#### Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.
- · We use the Expasy's ProtParam tool to determine the absorption coefficient of each protein.

#### Purification:

Two step purification of proteins expressed in Almost Living Cell-Free Expression System (ALiCE®):

- 1. In a first purification step, the protein is purified from the cleared cell lysate using StrepTag capture material. Eluate fractions are analyzed by SDS-PAGE.
- Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

## **Product Details** >80 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot. Purity: Endotoxin Level: Low Endotoxin less than 1 EU/mg (< 0.1 ng/mg) Grade: Crystallography grade **Target Details** MKNK2 Target: Alternative Name: MKNK2 (MKNK2 Products) Background: MAP kinase-interacting serine/threonine-protein kinase 2 (EC 2.7.11.1) (MAP kinase signalintegrating kinase 2) (MAPK signal-integrating kinase 2) (Mnk2), FUNCTION: Serine/threonineprotein kinase that phosphorylates SFPQ/PSF, HNRNPA1 and EIF4E. May play a role in the response to environmental stress and cytokines. Appears to regulate translation by phosphorylating EIF4E, thus increasing the affinity of this protein for the 7-methylguanosinecontaining mRNA cap. Required for mediating PP2A-inhibition-induced EIF4E phosphorylation. Triggers EIF4E shuttling from cytoplasm to nucleus. Isoform 1 displays a high basal kinase activity, but isoform 2 exhibits a very low kinase activity. Acts as a mediator of the suppressive effects of IFNgamma on hematopoiesis. Negative regulator for signals that control generation of arsenic trioxide As(2)O(3)-dependent apoptosis and anti-leukemic responses. Involved in anti-apoptotic signaling in response to serum withdrawal. {ECO:0000269|PubMed:11154262, ECO:0000269|PubMed:11463832, ECO:0000269|PubMed:12897141, ECO:0000269|PubMed:16111636, ECO:0000269|PubMed:17965020, ECO:0000269|PubMed:18299328, ECO:0000269|PubMed:20823271, ECO:0000269|PubMed:20927323, ECO:0000269|PubMed:21149447}. Molecular Weight: 51.9 kDa UniProt: Q9HBH9 Pathways: MAPK Signaling **Application Details**

Application Notes:

In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.

Comment:

ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce

even the most difficult-to-express proteins, including those that require post-translational modifications.

During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!


Restrictions:

For Research Use only

#### Handling

Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer. If you have a special request, please contact us.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	Unlimited (if stored properly)


#### **Images**



MAP Kinase Interacting serine/threonine Kinase 2 (MKNK2) (AA 1 – 465), gel filtration, Superose 6, fraction 12 - 13

Size-exclusion chromatography-High Pressure Liquid Chromatography

Image 1. SEC



MAP Kinase Interacting serine/threonine Kinase 2 (MKNK2) (AA 1 - 465), gel filtration, fraction 12 - 13

#### **Western Blotting**

Image 2. Quality Control Images: Western Blotting + SDS-PAGE