# antibodies .- online.com





### MYSM1 Protein (AA 1-828) (His tag)



**Image** 



Go to Product page

#### Overview

Quantity:	1 mg
Target:	MYSM1
Protein Characteristics:	AA 1-828
Origin:	Human
Source:	Insect Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This MYSM1 protein is labelled with His tag.
Application:	ELISA, Western Blotting (WB), Crystallization (Crys), SDS-PAGE (SDS)

#### **Product Details**

Sequence:

MAAEEADVDI EGDVVAAAGA QPGSGENTAS VLQKDHYLDS SWRTENGLIP WTLDNTISEE
NRAVIEKMLL EEEYYLSKKS QPEKVWLDQK EDDKKYMKSL QKTAKIMVHS PTKPASYSVK
WTIEEKELFE QGLAKFGRRW TKISKLIGSR TVLQVKSYAR QYFKNKVKCG LDKETPNQKT
GHNLQVKNED KGTKAWTPSC LRGRADPNLN AVKIEKLSDD EEVDITDEVD ELSSQTPQKN
SSSDLLLDFP NSKMHETNQG EFITSDSQEA LFSKSSRGCL QNEKQDETLS SSEITLWTEK
QSNGDKKSIE LNDQKFNELI KNCNKHDGRG IIVDARQLPS PEPCEIQKNL NDNEMLFHSC
QMVEESHEEE ELKPPEQEIE IDRNIIQEEE KQAIPEFFEG RQAKTPERYL KIRNYILDQW
EICKPKYLNK TSVRPGLKNC GDVNCIGRIH TYLELIGAIN FGCEQAVYNR PQTVDKVRIR
DRKDAVEAYQ LAQRLQSMRT RRRRVRDPWG NWCDAKDLEG QTFEHLSAEE LAKRREEEKG
RPVKSLKVPR PTKSSFDPFQ LIPCNFFSEE KQEPFQVKVA SEALLIMDLH AHVSMAEVIG
LLGGRYSEVD KVVEVCAAEP CNSLSTGLQC EMDPVSQTQA SETLAVRGFS VIGWYHSHPA
FDPNPSLRDI DTQAKYQSYF SRGGAKFIGM IVSPYNRNNP LPYSQITCLV ISEEISPDGS

YRLPYKFEVQ QMLEEPQWGL VFEKTRWIIE KYRLSHSSVP MDKIFRRDSD LTCLQKLLEC MRKTLSKVTN CFMAEEFLTE IENLFLSNYK SNQENGVTEE NCTKELLM

Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a special request, please contact us.

#### Characteristics:

- Made in Germany from design to production by highly experienced protein experts.
- Human MYSM1 Protein (raised in Insect Cells) purified by multi-step, protein-specific process to ensure crystallization grade.
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made to order protein and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).

When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered.

The concentration of our recombinant proteins is measured using the absorbance at 280nm.

The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.

The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

#### Purification:

Two step purification of proteins expressed in baculovirus infected SF9 insect cells:

- 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.
- Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

Purity:

>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.

Sterility:

0.22 µm filtered

Endotoxin Level:

Protein is endotoxin free

## Grade: Crystallography grade **Target Details** Target: MYSM1 Alternative Name MYSM1 (MYSM1 Products) Background: Metalloprotease that specifically deubiquitinates monoubiquitinated histone H2A, a specific tag for epigenetic transcriptional repression, thereby acting as a coactivator. Preferentially deubiquitinates monoubiquitinated H2A in hyperacetylated nucleosomes. Deubiquitination of histone H2A leads to facilitate the phosphorylation and dissociation of histone H1 from the nucleosome. Acts as a coactivator by participating in the initiation and elongation steps of androgen receptor (AR)-induced gene activation. {ECO:0000269|PubMed:17707232}. Molecular Weight: 96.0 kDa Including tag. UniProt: Q5VVJ2 Pathways: **Chromatin Binding Application Details** In addition to the applications listed above we expect the protein to work for functional studies **Application Notes:** as well. As the protein has not been tested for functional studies yet we cannot offer a gurantee though. Comment: In cases in which it is highly likely that the recombinant protein with the default tag will be insoluble our protein lab may suggest a higher molecular weight tag (e.g. GST-tag) instead to increase solubility. We will discuss all possible options with you in detail to assure that you receive your protein of interest. Restrictions: For Research Use only Handling Format: Liquid Buffer: 100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer. Handling Advice: Avoid repeated freeze-thaw cycles. Storage: -80 °C

**Product Details** 

Storage Comment:

Store at -80°C.

Expiry Date:

Unlimited (if stored properly)

**Images** 



**Image 1.** "Crystallography Grade" protein due to multi-step, protein-specific purification process