

# Datasheet for ABIN3094076 **NEK6 Protein (AA 1-313) (Strep Tag)**



### Overview

Quantity:	250 μg
Target:	NEK6
Protein Characteristics:	AA 1-313
Origin:	Human
Source:	Cell-free protein synthesis (CFPS)
Protein Type:	Recombinant
Purification tag / Conjugate:	This NEK6 protein is labelled with Strep Tag.
Application:	ELISA, Western Blotting (WB), SDS-PAGE (SDS)

Product Details	
Brand:	AliCE®
Sequence:	MAGQPGHMPH GGSSNNLCHT LGPVHPPDPQ RHPNTLSFRC SLADFQIEKK IGRGQFSEVY
	KATCLLDRKT VALKKVQIFE MMDAKARQDC VKEIGLLKQL NHPNIIKYLD SFIEDNELNI
	VLELADAGDL SQMIKYFKKQ KRLIPERTVW KYFVQLCSAV EHMHSRRVMH RDIKPANVFI
	TATGVVKLGD LGLGRFFSSE TTAAHSLVGT PYYMSPERIH ENGYNFKSDI WSLGCLLYEM
	AALQSPFYGD KMNLFSLCQK IEQCDYPPLP GEHYSEKLRE LVSMCICPDP HQRPDIGYVH
	QVAKQMHIWM SST
	Sequence without tag. The proposed Strep-Tag is based on experience s with the expression
	system, a different complexity of the protein could make another tag necessary. In case you
	have a special request, please contact us.
Characteristics:	Key Benefits:

- Made in Germany from design to production by highly experienced protein experts.
- · Protein expressed with ALiCE® and purified in one-step affinity chromatography
- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- · State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

#### Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require posttranslational modifications.
- During lysate production, the cell wall and other cellular components that are not required for
  protein production are removed, leaving only the protein production machinery and the
  mitochondria to drive the reaction. During our lysate completion steps, the additional
  components needed for protein production (amino acids, cofactors, etc.) are added to
  produce something that functions like a cell, but without the constraints of a living system all that's needed is the DNA that codes for the desired protein!

#### Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured against its specific reference buffer.
- · We use the Expasy's ProtParam tool to determine the absorption coefficient of each protein.

Purification:	One-step Strep-tag purification of proteins expressed in Almost Living Cell-Free Expression System (AliCE®).
Purity:	> 70-80 % as determined by SDS PAGE, Western Blot and analytical SEC (HPLC).
Grade:	custom-made
Target Details	
Target:	NEK6

## **Target Details**

Alternative Name:	NEK6 (NEK6 Products)
Background:	Serine/threonine-protein kinase Nek6 (EC 2.7.11.34) (Never in mitosis A-related kinase 6)
	(NimA-related protein kinase 6) (Protein kinase SID6-1512),FUNCTION: Protein kinase which
	plays an important role in mitotic cell cycle progression (PubMed:11516946,
	PubMed:14563848). Required for chromosome segregation at metaphase-anaphase transition
	robust mitotic spindle formation and cytokinesis (PubMed:19414596). Phosphorylates ATF4,
	CIR1, PTN, RAD26L, RBBP6, RPS7, RPS6KB1, TRIP4, STAT3 and histones H1 and H3
	(PubMed:12054534, PubMed:20873783). Phosphorylates KIF11 to promote mitotic spindle
	formation (PubMed:19001501). Involved in G2/M phase cell cycle arrest induced by DNA
	damage (PubMed:18728393). Inhibition of activity results in apoptosis. May contribute to
	tumorigenesis by suppressing p53/TP53-induced cancer cell senescence (PubMed:21099361)
	Phosphorylates EML4 at 'Ser-144', promoting its dissociation from microtubules during mitosis
	which is required for efficient chromosome congression (PubMed:31409757).
	{ECO:0000269 PubMed:11516946, ECO:0000269 PubMed:12054534,
	ECO:0000269 PubMed:14563848, ECO:0000269 PubMed:18728393,
	ECO:0000269 PubMed:19001501, ECO:0000269 PubMed:19414596,
	ECO:0000269 PubMed:20873783, ECO:0000269 PubMed:21099361,
	ECO:0000269 PubMed:31409757}.
Molecular Weight:	35.7 kDa
UniProt:	Q9HC98
Application Details	
Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies
	as well. As the protein has not been tested for functional studies yet we cannot offer a
	guarantee though.
Comment:	ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from
Comment:	
Comment:	
Comment:	Nicotiana tabacum c.v This contains all the protein expression machinery needed to produce
Comment:	Nicotiana tabacum c.v This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational
Comment:	Nicotiana tabacum c.v This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.
Comment:	Nicotiana tabacum c.v This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.  During lysate production, the cell wall and other cellular components that are not required for
Comment:	Nicotiana tabacum c.v This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.  During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the

## **Application Details**

	needed is the DNA that codes for the desired protein!
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer.  Standard Storage Buffer: PBS pH 7.4, 10 % Glycerol <b>Might differ depending on protein.</b>
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	12 months