# antibodies .- online.com





# TRKA Protein (AA 33-423) (His tag)



# **Image**



#### Overview

Quantity:	1 mg
Target:	TRKA (NTRK1)
Protein Characteristics:	AA 33-423
Origin:	Human
Source:	Insect Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This TRKA protein is labelled with His tag.
Application:	Western Blotting (WB), SDS-PAGE (SDS), ELISA, Crystallization (Crys)

#### **Product Details**

### Sequence:

AAPCPDACCP HGSSGLRCTR DGALDSLHHL PGAENLTELY IENQQHLQHL ELRDLRGLGE
LRNLTIVKSG LRFVAPDAFH FTPRLSRLNL SFNALESLSW KTVQGLSLQE LVLSGNPLHC
SCALRWLQRW EEEGLGGVPE QKLQCHGQGP LAHMPNASCG VPTLKVQVPN ASVDVGDDVL
LRCQVEGRGL EQAGWILTEL EQSATVMKSG GLPSLGLTLA NVTSDLNRKN VTCWAENDVG
RAEVSVQVNV SFPASVQLHT AVEMHHWCIP FSVDGQPAPS LRWLFNGSVL NETSFIFTEF
LEPAANETVR HGCLRLNQPT HVNNGNYTLL AANPFGQASA SIMAAFMDNP FEFNPEDPIP
VSFSPVDTNS TSGDPVEKKD ETPFGVSVAV G

Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a special request, please contact us.

#### Characteristics:

- Made in Germany from design to production by highly experienced protein experts.
- Human NTRK1 Protein (raised in Insect Cells) purified by multi-step, protein-specific process to ensure crystallization grade.

• State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made to order protein and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).

When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered. The concentration of our recombinant proteins is measured using the absorbance at 280nm. The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.

The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

Purification:

Two step purification of proteins expressed in baculovirus infected SF9 insect cells:

- In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.
- Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

 Purity:
 >95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.

 Sterility:
 0.22 μm filtered

 Endotoxin Level:
 Protein is endotoxin free.

 Grade:
 Crystallography grade

#### **Target Details**

Target: TRKA (NTRK1)

Alternative Name: NTRK1 (NTRK1 Products)

Background:

Receptor tyrosine kinase involved in the development and the maturation of the central and peripheral nervous systems through regulation of proliferation, differentiation and survival of sympathetic and nervous neurons. High affinity receptor for NGF which is its primary ligand, it can also bind and be activated by NTF3/neurotrophin-3. However, NTF3 only supports axonal extension through NTRK1 but has no effect on neuron survival. Upon dimeric NGF ligandbinding, undergoes homodimerization, autophosphorylation and activation. Recruits, phosphorylates and/or activates several downstream effectors including SHC1, FRS2, SH2B1, SH2B2 and PLCG1 that regulate distinct overlapping signaling cascades driving cell survival and differentiation. Through SHC1 and FRS2 activates a GRB2-Ras-MAPK cascade that regulates cell differentiation and survival. Through PLCG1 controls NF-Kappa-B activation and the transcription of genes involved in cell survival. Through SHC1 and SH2B1 controls a Ras-PI3 kinase-AKT1 signaling cascade that is also regulating survival. In absence of ligand and activation, may promote cell death, making the survival of neurons dependent on trophic factors., Isoform TrkA-III is resistant to NGF, constitutively activates AKT1 and NF-kappa-B and is unable to activate the Ras-MAPK signaling cascade. Antagonizes the anti-proliferative NGF-NTRK1 signaling that promotes neuronal precursors differentiation. Isoform TrkA-III promotes angiogenesis and has oncogenic activity when overexpressed.

Molecular Weight:

43.4 kDa Including tag.

UniProt:

P04629

Pathways:

RTK Signaling, Neurotrophin Signaling Pathway, cAMP Metabolic Process

# Application Details

Application Notes:

In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a gurantee though.

Comment:

In cases in which it is highly likely that the recombinant protein with the default tag will be insoluble our protein lab may suggest a higher molecular weight tag (e.g. GST-tag) instead to increase solubility. We will discuss all possible options with you in detail to assure that you receive your protein of interest.

Restrictions:

For Research Use only

## Handling


Format:

Liquid

# Handling

Buffer:	100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	Unlimited (if stored properly)

# **Images**



**Image 1.** "Crystallography Grade" protein due to multi-step, protein-specific purification process