

Datasheet for ABIN3094770

PRDM1 Protein (AA 1-825) (Strep Tag)



()	ve	rvi	6	W
\sim	v C	1 V I	\sim	v v

Quantity:	250 μg
Target:	PRDM1
Protein Characteristics:	AA 1-825
Origin:	Human
Source:	Cell-free protein synthesis (CFPS)
Protein Type:	Recombinant
Purification tag / Conjugate:	This PRDM1 protein is labelled with Strep Tag.
Application:	Western Blotting (WB), SDS-PAGE (SDS), ELISA

Product Details				
Brand:	AliCE®			
Sequence:	MLDICLEKRV GTTLAAPKCN SSTVRFQGLA EGTKGTMKMD MEDADMTLWT EAEFEEKCTY			
	IVNDHPWDSG ADGGTSVQAE ASLPRNLLFK YATNSEEVIG VMSKEYIPKG TRFGPLIGEI			
	YTNDTVPKNA NRKYFWRIYS RGELHHFIDG FNEEKSNWMR YVNPAHSPRE QNLAACQNGM			
	NIYFYTIKPI PANQELLVWY CRDFAERLHY PYPGELTMMN LTQTQSSLKQ PSTEKNELCP			
	KNVPKREYSV KEILKLDSNP SKGKDLYRSN ISPLTSEKDL DDFRRRGSPE MPFYPRVVYP			
	IRAPLPEDFL KASLAYGIER PTYITRSPIP SSTTPSPSAR SSPDQSLKSS SPHSSPGNTV			
	SPVGPGSQEH RDSYAYLNAS YGTEGLGSYP GYAPLPHLPP AFIPSYNAHY PKFLLPPYGM			
	NCNGLSAVSS MNGINNFGLF PRLCPVYSNL LGGGSLPHPM LNPTSLPSSL PSDGARRLLQ			
	PEHPREVLVP APHSAFSFTG AAASMKDKAC SPTSGSPTAG TAATAEHVVQ PKATSAAMAA			
	PSSDEAMNLI KNKRNMTGYK TLPYPLKKQN GKIKYECNVC AKTFGQLSNL KVHLRVHSGE			
	RPFKCQTCNK GFTQLAHLQK HYLVHTGEKP HECQVCHKRF SSTSNLKTHL RLHSGEKPYQ			

CKVCPAKFTQ FVHLKLHKRL HTRERPHKCS QCHKNYIHLC SLKVHLKGNC AAAPAPGLPL EDLTRINEEI EKFDISDNAD RLEDVEDDIS VISVVEKEIL AVVRKEKEET GLKVSLQRNM GNGLLSSGCS LYESSDLPLM KLPPSNPLPL VPVKVKQETV EPMDP

Sequence without tag. The proposed Strep-Tag is based on experience s with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.

Characteristics:

Key Benefits:

- Made in Germany from design to production by highly experienced protein experts.
- Protein expressed with ALiCE® and purified in one-step affinity chromatography
- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require posttranslational modifications.
- During lysate production, the cell wall and other cellular components that are not required for
 protein production are removed, leaving only the protein production machinery and the
 mitochondria to drive the reaction. During our lysate completion steps, the additional
 components needed for protein production (amino acids, cofactors, etc.) are added to
 produce something that functions like a cell, but without the constraints of a living system all that's needed is the DNA that codes for the desired protein!

Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured against its specific reference buffer.
- We use the Expasy's ProtParam tool to determine the absorption coefficient of each protein.

Purification:

One-step Strep-tag purification of proteins expressed in Almost Living Cell-Free Expression

Product Details

Product Details		
	System (AliCE®).	
Purity:	> 70-80 % as determined by SDS PAGE, Western Blot and analytical SEC (HPLC).	
Grade:	custom-made	
Target Details		
Target:	PRDM1	
Alternative Name:	PRDM1 (PRDM1 Products)	
Background:	PR domain zinc finger protein 1 (EC 2.1.1) (BLIMP-1) (Beta-interferon gene positive regulatory	
	domain I-binding factor) (PR domain-containing protein 1) (Positive regulatory domain I-binding	
	factor 1) (PRDI-BF1) (PRDI-binding factor 1),FUNCTION: Transcription factor that mediates a	
	transcriptional program in various innate and adaptive immune tissue-resident lymphocyte T	
	cell types such as tissue-resident memory T (Trm), natural killer (trNK) and natural killer T (NKT)	
	cells and negatively regulates gene expression of proteins that promote the egress of tissue-	
	resident T-cell populations from non-lymphoid organs. Plays a role in the development,	
	retention and long-term establishment of adaptive and innate tissue-resident lymphocyte T cell	
	types in non-lymphoid organs, such as the skin and gut, but also in other nonbarrier tissues like	
	liver and kidney, and therefore may provide immediate immunological protection against	
	reactivating infections or viral reinfection (By similarity). Binds specifically to the PRDI element	
	in the promoter of the beta-interferon gene (PubMed:1851123). Drives the maturation of B-	
	lymphocytes into Ig secreting cells (PubMed:12626569). Associates with the transcriptional	
	repressor ZNF683 to chromatin at gene promoter regions (By similarity). Binds to the promoter	
	and acts as a transcriptional repressor of IRF8, thereby promotes transcription of osteoclast	
	differentiation factors such as NFATC1 and EEIG1 (By similarity).	
	{ECO:0000250 UniProtKB:Q60636, ECO:0000269 PubMed:12626569,	
	ECO:0000269 PubMed:1851123}.	
Molecular Weight:	91.8 kDa	
UniProt:	075626	
Pathways:	Regulation of Muscle Cell Differentiation	
Application Details		
Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies	
	as well. As the protein has not been tested for functional studies yet we cannot offer a	

Application Details

Storage Comment:

Expiry Date:

Application Details		
	guarantee though.	
Comment:	ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from	
	Nicotiana tabacum c.v This contains all the protein expression machinery needed to produce	
	even the most difficult-to-express proteins, including those that require post-translational	
	modifications.	
	During lysate production, the cell wall and other cellular components that are not required for	
	protein production are removed, leaving only the protein production machinery and the	
	mitochondria to drive the reaction. During our lysate completion steps, the additional	
	components needed for protein production (amino acids, cofactors, etc.) are added to produce	
	something that functions like a cell, but without the constraints of a living system - all that's	
	needed is the DNA that codes for the desired protein!	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Buffer:	The buffer composition is at the discretion of the manufacturer.	
	Standard Storage Buffer: PBS pH 7.4, 10 % Glycerol Might differ depending on protein.	
Handling Advice:	Avoid repeated freeze-thaw cycles.	
Storage:	-80 °C	

Store at -80°C.

12 months