

Datasheet for ABIN3095071

## RNF146 Protein (AA 1-359) (Strep Tag)



[Go to Product page](#)

### Overview

Quantity:	250 µg
Target:	RNF146
Protein Characteristics:	AA 1-359
Origin:	Human
Source:	Cell-free protein synthesis (CFPS)
Protein Type:	Recombinant
Purification tag / Conjugate:	This RNF146 protein is labelled with Strep Tag.
Application:	ELISA, Western Blotting (WB), SDS-PAGE (SDS)

### Product Details

Brand:	AliCE®
Sequence:	<p>MMAGCGEIDH SINMLPTNRK ANESCSNTAP SLTVPECAIC LQTCVHPVSL PCKHVFCYLC  VKGASWLGKR CALCRQEIPE DFLDKPTLLS PEELKAASRG NGEYAWYYEG RNGWWQYDER  TSRELEDAFS KGKKNTEMLI AGFLYVADLE NMVQYRRNEH GRRRKIKRDI IDIPKKGVAG  LRLDCDANTV NLARESSADG ADSVSAQSGA SVQPLVSSVR PLTSVDGQLT SPATPSPDAS  TSLEDSFAHL QLSGDNTAER SHRGEGEEDH ESPSSGRVPA PDTSIEETES DASDSEDVS  AVVAQHSLTQ QRLLVSNANQ TVPDRSDRSR TDRSVAGGGT VSVSVRSRRP DGQCTVTEV</p> <p><b>Sequence without tag. The proposed Strep-Tag is based on experience s with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.</b></p>
Characteristics:	Key Benefits:

## Product Details

---

- Made in Germany - from design to production - by highly experienced protein experts.
- Protein expressed with ALiCE® and purified in one-step affinity chromatography
- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

### Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from *Nicotiana tabacum* c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.
- During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

### Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured against its specific reference buffer.
- We use the Expasy's ProtParam tool to determine the absorption coefficient of each protein.

Purification:	One-step Strep-tag purification of proteins expressed in Almost Living Cell-Free Expression System (ALiCE®).
Purity:	> 70-80 % as determined by SDS PAGE, Western Blot and analytical SEC (HPLC).
Grade:	custom-made

## Target Details

---

Target:	RNF146
---------	--------

---

## Target Details

Alternative Name: RNF146 ([RNF146 Products](#))

Background: E3 ubiquitin-protein ligase RNF146 (EC 2.3.2.27) (Dactylidin) (Iduna) (RING finger protein 146) (RING-type E3 ubiquitin transferase RNF146),FUNCTION: E3 ubiquitin-protein ligase that specifically binds poly-ADP-ribosylated (PARsylated) proteins and mediates their ubiquitination and subsequent degradation (PubMed:21478859, PubMed:21799911, PubMed:22267412). May regulate many important biological processes, such as cell survival and DNA damage response (PubMed:21825151, PubMed:22267412). Acts as an activator of the Wnt signaling pathway by mediating the ubiquitination of PARsylated AXIN1 and AXIN2, 2 key components of the beta-catenin destruction complex (PubMed:21478859, PubMed:21799911). Acts in cooperation with tankyrase proteins (TNKS and TNKS2), which mediate PARsylation of target proteins AXIN1, AXIN2, BLZF1, CASC3, TNKS and TNKS2 (PubMed:21799911). Recognizes and binds tankyrase-dependent PARsylated proteins via its WWE domain and mediates their ubiquitination, leading to their degradation (PubMed:21799911). Different ubiquitin linkage types have been observed: TNKS2 undergoes ubiquitination at 'Lys-48' and 'Lys-63', while AXIN1 is only ubiquitinated at 'Lys-48' (PubMed:21799911). May regulate TNKS and TNKS2 subcellular location, preventing aggregation at a centrosomal location (PubMed:21799911). Neuroprotective protein (PubMed:21602803). Protects the brain against N-methyl-D-aspartate (NMDA) receptor-mediated glutamate excitotoxicity and ischemia, by interfering with PAR-induced cell death, called parthanatos (By similarity). Prevents nuclear translocation of AIFM1 in a PAR-binding dependent manner (By similarity). Does not affect PARP1 activation (By similarity). Protects against cell death induced by DNA damaging agents, such as N-methyl-N-nitro-N-nitrosoguanidine (MNNG) and rescues cells from G1 arrest (By similarity). Promotes cell survival after gamma-irradiation (PubMed:21825151). Facilitates DNA repair (PubMed:21825151). {ECO:0000250|UniProtKB:Q9CZW6, ECO:0000269|PubMed:21478859, ECO:0000269|PubMed:21602803, ECO:0000269|PubMed:21799911, ECO:0000269|PubMed:21825151, ECO:0000269|PubMed:22267412}.

Molecular Weight: 39.0 kDa

UniProt: [Q9NTX7](#)

## Application Details

Application Notes: In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.

Comment: ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from

Application Details

Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.

During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

Restrictions: For Research Use only

Handling

Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer. Standard Storage Buffer: PBS pH 7.4, 10 % Glycerol <b>Might differ depending on protein.</b>
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	12 months