

Datasheet for ABIN3095428  
**SETD7 Protein (AA 1-366) (Strep Tag)**



[Go to Product page](#)

1 Image

Overview

Quantity:	1 mg
Target:	SETD7
Protein Characteristics:	AA 1-366
Origin:	Human
Source:	Tobacco (Nicotiana tabacum)
Protein Type:	Recombinant
Purification tag / Conjugate:	This SETD7 protein is labelled with Strep Tag.
Application:	Western Blotting (WB), SDS-PAGE (SDS), ELISA

Product Details

Sequence: MDSDDDEMVEE AVEGHLDDDG LPHGFCTVTY SSTDRFEGNF VHGEKNGRGK FFFFDGSTLE  
GYYVDDALQG QGVYTYEDGG VLQGTYYVDGE LNGPAQEYDT DGRLLFKGQY KDNIRHGVCW  
IYYPDGGSLV GEVNEGEMT GEKIAYVYPD ERTALYGKFI DGEMIEGKLA TLMSTEEGRP  
HFELMPGNSV YHFDKSTSSC ISTNALLPDP YESERVYVAE SLISSAGEGL FSKVAVGPNT  
VMSFYNGVRI THQEVDSDRW ALNGNTLSLD EETVIDVPEP YNHVSKYCAS LGHKANHSFT  
PNCIYDMFVH PRFGPIKCIR TLRAVEADEE LTVAYGYDHS PPGKSGPEAP EWYQVELKAF  
QATQQK

**Sequence without tag. The proposed Strep-Tag is based on experience s with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.**

Characteristics: Key Benefits:

- Made in Germany - from design to production - by highly experienced protein experts.
- Protein expressed with ALiCE® and purified by multi-step, protein-specific process to ensure correct folding and modification.
- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

### Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from *Nicotiana tabacum* c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.
- During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

### Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.
- We use the Exspasy's ProtParam tool to determine the absorption coefficient of each protein.

---

### Purification:

Two step purification of proteins expressed in Almost Living Cell-Free Expression System (ALiCE®):

1. In a first purification step, the protein is purified from the cleared cell lysate using StrepTag capture material. Eluate fractions are analyzed by SDS-PAGE.
2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

## Product Details

---

Purity:	>80 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.
Endotoxin Level:	Low Endotoxin less than 1 EU/mg (< 0.1 ng/mg)
Grade:	Crystallography grade

## Target Details

---

Target:	SETD7
Alternative Name:	SETD7 ( <a href="#">SETD7 Products</a> )
Background:	<p>Histone-lysine N-methyltransferase SETD7 (EC 2.1.1.364) (Histone H3-K4 methyltransferase SETD7) (H3-K4-HMTase SETD7) (Lysine N-methyltransferase 7) (SET domain-containing protein 7) (SET7/9),FUNCTION: Histone methyltransferase that specifically monomethylates 'Lys-4' of histone H3 (PubMed:11779497, PubMed:11850410, PubMed:12588998, PubMed:12540855, PubMed:16141209). H3 'Lys-4' methylation represents a specific tag for epigenetic transcriptional activation (PubMed:12588998, PubMed:12540855, PubMed:16141209). Plays a central role in the transcriptional activation of genes such as collagenase or insulin (PubMed:16141209, PubMed:12588998). Recruited by IPF1/PDX-1 to the insulin promoter, leading to activate transcription (PubMed:16141209). Has also methyltransferase activity toward non-histone proteins such as CGAS, p53/TP53, TAF10, and possibly TAF7 by recognizing and binding the [KR]-[STA]-K in substrate proteins (PubMed:15099517, PubMed:35210392, PubMed:15525938, PubMed:16415881). Monomethylates 'Lys-189' of TAF10, leading to increase the affinity of TAF10 for RNA polymerase II (PubMed:15099517, PubMed:16415881). Monomethylates 'Lys-372' of p53/TP53, stabilizing p53/TP53 and increasing p53/TP53-mediated transcriptional activation (PubMed:17108971, PubMed:15525938, PubMed:16415881). Monomethylates 'Lys-491' of CGAS, promoting interaction between SGF29 and CGAS (By similarity). {ECO:0000250 UniProtKB:Q8VHL1, ECO:0000269 PubMed:11779497, ECO:0000269 PubMed:11850410, ECO:0000269 PubMed:12540855, ECO:0000269 PubMed:12588998, ECO:0000269 PubMed:15099517, ECO:0000269 PubMed:15525938, ECO:0000269 PubMed:16141209, ECO:0000269 PubMed:16415881, ECO:0000269 PubMed:17108971, ECO:0000269 PubMed:35210392}.</p>
Molecular Weight:	40.7 kDa
UniProt:	<a href="#">Q8WTS6</a>

## Application Details

---

**Application Notes:** In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.

---

**Comment:** ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from *Nicotiana tabacum* c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.

During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

---

**Restrictions:** For Research Use only

---

## Handling

---

**Format:** Liquid

---

**Buffer:** The buffer composition is at the discretion of the manufacturer. If you have a special request, please contact us.

---

**Handling Advice:** Avoid repeated freeze-thaw cycles.

---

**Storage:** -80 °C

---

**Storage Comment:** Store at -80°C.

---

**Expiry Date:** Unlimited (if stored properly)

---



**Image 1.** „Crystallography Grade“ protein due to multi-step, protein-specific purification process