antibodies -online.com





SPPL2A Protein (AA 26-172) (His tag)



Image



Go to Product page

()	11/	IN	/ie	A .
	/ // 	۱ ات	/ (−	' \/\/

Quantity:	1 mg	
Target:	SPPL2A	
Protein Characteristics:	AA 26-172	
Origin:	Human	
Source:	Escherichia coli (E. coli)	
Protein Type:	Recombinant	
Purification tag / Conjugate:	This SPPL2A protein is labelled with His tag.	
Application:	ELISA, Western Blotting (WB), Crystallization (Crys), SDS-PAGE (SDS)	
Product Details		
Sequence:	QEAILHASGN GTTKDYCMLY NPYWTALPST LENATSISLM NLTSTPLCNL SDIPPVGIKS	
	KAVVVPWGSC HFLEKARIAQ KGGAEAMLVV NNSVLFPPSG NRSEFPDVKI LIAFISYKDF	
	RDMNQTLGDN ITVKMYSPSW PNFDYTM	
	Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a	
	special request, please contact us.	
Characteristics:	 Made in Germany - from design to production - by highly experienced protein experts. Human SPPL2A Protein (raised in E. Coli) purified by multi-step, protein-specific process to ensure crystallization grade. State-of-the-art algorithm used for plasmid design (Gene synthesis). 	
	This protein is a made to order protein and will be made for the first time for your order. Our	
	experts in the lab will ensure that you receive a correctly folded protein.	
	The big advantage of ordering our made-to-order proteins in comparison to ordering custom	

made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).

When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered.

The concentration of our recombinant proteins is measured using the absorbance at 280nm.

The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.

The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

Purification:

Two step purification of proteins expressed in bacterial culture:

- 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.
- 2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

Purity:

>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.

Sterility:

0.22 µm filtered

Endotoxin Level:

Endotoxin has not been removed. Please contact us if you require endotoxin removal.

Grade:

Crystallography grade

Target Details

Target:	SPPL2A	
Alternative Name:	SPPL2A (SPPL2A Products)	
Background:	Intramembrane-cleaving aspartic protease (I-CLiP) that cleaves type II membrane signal	
	peptides in the hydrophobic plane of the membrane. Functions in FASLG, ITM2B and TNF	
	processing (PubMed:16829952, PubMed:16829951, PubMed:17557115, PubMed:17965014).	
	Catalyzes the intramembrane cleavage of the anchored fragment of shed TNF-alpha (TNF),	

which promotes the release of the intracellular domain (ICD) for signaling to the nucleus (PubMed:16829952). Also responsible for the intramembrane cleavage of Fas antigen ligand FASLG, which promotes the release of the intracellular FasL domain (FasL ICD) (PubMed:17557115). May play a role in the regulation of innate and adaptive immunity (PubMed:16829952). Catalyzes the intramembrane cleavage of the simian foamy virus envelope glycoprotein gp130 independently of prior ectodomain shedding by furin or furin-like proprotein convertase (PC)-mediated cleavage proteolysis (PubMed:23132852). {ECO:0000269|PubMed:16829951, ECO:0000269|PubMed:16829952, ECO:0000269|PubMed:17557115, ECO:0000269|PubMed:17965014, ECO:0000269|PubMed:23132852}.

Molecular Weight:

17.2 kDa Including tag.

UniProt:

Q8TCT8

Application Details

Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies
	as well. As the protein has not been tested for functional studies yet we cannot offer a gurantee
	though.

Comment:

In cases in which it is highly likely that the recombinant protein with the default tag will be insoluble our protein lab may suggest a higher molecular weight tag (e.g. GST-tag) instead to increase solubility. We will discuss all possible options with you in detail to assure that you receive your protein of interest.

Restrictions:

For Research Use only

Handling

Format:	Liquid	
Buffer:	100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer.	
Handling Advice:	Avoid repeated freeze-thaw cycles.	
Storage:	-80 °C	
Storage Comment:	Store at -80°C.	
Expiry Date:	Unlimited (if stored properly)	



Image 1. "Crystallography Grade" protein due to multi-step, protein-specific purification process