

Datasheet for ABIN3095776

TARDBP Protein (AA 1-414) (Strep Tag)



Overview

Quantity:	250 μg
Target:	TARDBP
Protein Characteristics:	AA 1-414
Origin:	Human
Source:	Cell-free protein synthesis (CFPS)
Protein Type:	Recombinant
Purification tag / Conjugate:	This TARDBP protein is labelled with Strep Tag.
Application:	SDS-PAGE (SDS), Western Blotting (WB), ELISA

Product Details	
Brand:	AliCE®
Sequence:	MSEYIRVTED ENDEPIEIPS EDDGTVLLST VTAQFPGACG LRYRNPVSQC MRGVRLVEGI
	LHAPDAGWGN LVYVVNYPKD NKRKMDETDA SSAVKVKRAV QKTSDLIVLG LPWKTTEQDL
	KEYFSTFGEV LMVQVKKDLK TGHSKGFGFV RFTEYETQVK VMSQRHMIDG RWCDCKLPNS
	KQSQDEPLRS RKVFVGRCTE DMTEDELREF FSQYGDVMDV FIPKPFRAFA FVTFADDQIA
	QSLCGEDLII KGISVHISNA EPKHNSNRQL ERSGRFGGNP GGFGNQGGFG NSRGGGAGLG
	NNQGSNMGGG MNFGAFSINP AMMAAAQAAL QSSWGMMGML ASQQNQSGPS
	GNNQNQGNMQ REPNQAFGSG NNSYSGSNSG AAIGWGSASN AGSGSGFNGG FGSSMDSKSS
	GWGM
	Sequence without tag. The proposed Strep-Tag is based on experience s with the expression
	system, a different complexity of the protein could make another tag necessary. In case you
	have a special request, please contact us.

Characteristics:

Key Benefits:

- Made in Germany from design to production by highly experienced protein experts.
- · Protein expressed with ALiCE® and purified in one-step affinity chromatography
- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require posttranslational modifications.
- During lysate production, the cell wall and other cellular components that are not required for
 protein production are removed, leaving only the protein production machinery and the
 mitochondria to drive the reaction. During our lysate completion steps, the additional
 components needed for protein production (amino acids, cofactors, etc.) are added to
 produce something that functions like a cell, but without the constraints of a living system all that's needed is the DNA that codes for the desired protein!

Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured against its specific reference buffer.
- We use the Expasy's ProtParam tool to determine the absorption coefficient of each protein.

Purification:	One-step Strep-tag purification of proteins expressed in Almost Living Cell-Free Expression System (AliCE®).
Purity:	> 70-80 % as determined by SDS PAGE, Western Blot and analytical SEC (HPLC).
Grade:	custom-made

Target Details

Target:	TARDBP
Alternative Name:	TARDBP (TARDBP Products)
Background:	TAR DNA-binding protein 43 (TDP-43), FUNCTION: RNA-binding protein that is involved in
	various steps of RNA biogenesis and processing (PubMed:23519609). Preferentially binds, via
	its two RNA recognition motifs RRM1 and RRM2, to GU-repeats on RNA molecules
	predominantly localized within long introns and in the 3'UTR of mRNAs (PubMed:23519609,
	PubMed:24240615, PubMed:24464995). In turn, regulates the splicing of many non-coding and
	protein-coding RNAs including proteins involved in neuronal survival, as well as mRNAs that
	encode proteins relevant for neurodegenerative diseases (PubMed:21358640,
	PubMed:29438978). Plays a role in maintaining mitochondrial homeostasis by regulating the
	processing of mitochondrial transcripts (PubMed:28794432). Regulates also mRNA stability by
	recruiting CNOT7/CAF1 deadenylase on mRNA 3'UTR leading to poly(A) tail deadenylation and
	thus shortening (PubMed:30520513). In response to oxidative insult, associates with stalled
	ribosomes localized to stress granules (SGs) and contributes to cell survival
	(PubMed:23398327, PubMed:19765185). Participates also in the normal skeletal muscle
	formation and regeneration, forming cytoplasmic myo-granules and binding mRNAs that
	encode sarcomeric proteins (PubMed:30464263). Plays a role in the maintenance of the
	circadian clock periodicity via stabilization of the CRY1 and CRY2 proteins in a FBXL3-
	dependent manner (PubMed:27123980). Negatively regulates the expression of CDK6
	(PubMed:19760257). Regulates the expression of HDAC6, ATG7 and VCP in a PPIA/CYPA-
	dependent manner (PubMed:25678563). {ECO:0000269 PubMed:11285240,
	ECO:0000269 PubMed:17481916, ECO:0000269 PubMed:19760257,
	ECO:0000269 PubMed:19765185, ECO:0000269 PubMed:21358640,
	ECO:0000269 PubMed:23398327, ECO:0000269 PubMed:23519609,
	ECO:0000269 PubMed:24240615, ECO:0000269 PubMed:24464995,
	ECO:0000269 PubMed:25678563, ECO:0000269 PubMed:27123980,
	ECO:0000269 PubMed:28794432, ECO:0000269 PubMed:29438978,
	ECO:0000269 PubMed:30464263, ECO:0000269 PubMed:30520513}.
Molecular Weight:	44.7 kDa
UniProt:	Q13148
Pathways:	Positive Regulation of Peptide Hormone Secretion

Application Details

Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.
Comment:	ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from
	Nicotiana tabacum c.v This contains all the protein expression machinery needed to produce
	even the most difficult-to-express proteins, including those that require post-translational
	modifications.
	During lysate production, the cell wall and other cellular components that are not required for
	protein production are removed, leaving only the protein production machinery and the
	mitochondria to drive the reaction. During our lysate completion steps, the additional
	components needed for protein production (amino acids, cofactors, etc.) are added to produce
	something that functions like a cell, but without the constraints of a living system - all that's
	needed is the DNA that codes for the desired protein!
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer.
	Standard Storage Buffer: PBS pH 7.4, 10 % Glycerol Might differ depending on protein.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	12 months