# antibodies .- online.com





# TCF7L1 Protein (AA 1-588) (Strep Tag)



## **Image**



Go to Product page

#### Overview

Quantity:	1 mg
Target:	TCF7L1
Protein Characteristics:	AA 1-588
Origin:	Human
Source:	Tobacco (Nicotiana tabacum)
Protein Type:	Recombinant
Purification tag / Conjugate:	This TCF7L1 protein is labelled with Strep Tag.
Application:	ELISA, Western Blotting (WB), SDS-PAGE (SDS)

## **Product Details**

Sequence:

MPQLGGGGG GGGGSGGGG SSAGAAGGGD DLGANDELIP FQDEGGEEQE PSSDSASAQR DLDEVKSSLV NESENQSSSS DSEAERRPQP VRDTFQKPRD YFAEVRRPQD SAFFKGPPYP GYPFLMIPDL SSPYLSNGPL SPGGARTYLQ MKWPLLDVPS SATVKDTRSP SPAHLSNKVP VVQHPHHMHP LTPLITYSND HFSPGSPPTH LSPEIDPKTG IPRPPHPSEL SPYYPLSPGA VGQIPHPLGW LVPQQGQPMY SLPPGGFRHP YPALAMNASM SSLVSSRFSP HMVAPAHPGL PTSGIPHPAI VSPIVKQEPA PPSLSPAVSV KSPVTVKKEE EKKPHVKKPL NAFMLYMKEM RAKVVAECTL KESAAINQIL GRKWHNLSRE EQAKYYELAR KERQLHSQLY PTWSARDNYG KKKKRKREKQ LSQTQSQQQV QEAEGALASK SKKPCVQYLP PEKPCDSPAS SHGSMLDSPA TPSAALASPA APAATHSEQA QPLSLTTKPE TRAQLALHSA AFLSAKAAAS SSGQMGSQPP LLSRPLPLGS MPTALLASPP SFPATLHAHQ ALPVLQAQPL SLVTKSAH

Sequence without tag. The proposed Strep-Tag is based on experience s with the expression system, a different complexity of the protein could make another tag necessary. In case you

#### have a special request, please contact us.

#### Characteristics:

## Key Benefits:

- Made in Germany from design to production by highly experienced protein experts.
- Protein expressed with ALiCE® and purified by multi-step, protein-specific process to ensure correct folding and modification.
- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

## Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require posttranslational modifications.
- During lysate production, the cell wall and other cellular components that are not required for
  protein production are removed, leaving only the protein production machinery and the
  mitochondria to drive the reaction. During our lysate completion steps, the additional
  components needed for protein production (amino acids, cofactors, etc.) are added to
  produce something that functions like a cell, but without the constraints of a living system all that's needed is the DNA that codes for the desired protein!

## Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.
- We use the Expasy's ProtParam tool to determine the absorption coefficient of each protein.

## Purification:

Two step purification of proteins expressed in Almost Living Cell-Free Expression System (ALiCE®):

1. In a first purification step, the protein is purified from the cleared cell lysate using StrepTag capture material. Eluate fractions are analyzed by SDS-PAGE.

	<ol><li>Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.</li></ol>
Purity:	>80 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.
Endotoxin Level:	Low Endotoxin less than 1 EU/mg (< 0.1 ng/mg)
Grade:	Crystallography grade
Target Details	
Target:	TCF7L1
Alternative Name:	TCF7L1 (TCF7L1 Products)
Background:	Transcription factor 7-like 1 (HMG box transcription factor 3) (TCF-3),FUNCTION: Participates in the Wnt signaling pathway. Binds to DNA and acts as a repressor in the absence of CTNNB1, and as an activator in its presence. Necessary for the terminal differentiation of epidermal cells, the formation of keratohyalin granules and the development of the barrier function of the epidermis (By similarity). Down-regulates NQO1, leading to increased mitomycin c resistance. {ECO:0000250}.
Molecular Weight:	62.6 kDa
UniProt:	Q9HCS4
Pathways:	WNT Signaling, Stem Cell Maintenance
Application Details	
Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.
Comment:	ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.  During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce

## **Application Details**

**Images** 

	something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer. If you have a special request, please contact us.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	Unlimited (if stored properly)



Image 1. "Crystallography Grade" protein due to multi-step, protein-specific purification process