# antibodies -online.com





# TRPV4 Protein (AA 712-871) (His tag)



Go to Product page

$\sim$	
( )\/△	rview
$\circ$	1 410 44

0.00000	
Quantity:	1 mg
Target:	TRPV4
Protein Characteristics:	AA 712-871
Origin:	Human
Source:	Escherichia coli (E. coli)
Protein Type:	Recombinant
Purification tag / Conjugate:	This TRPV4 protein is labelled with His tag.
Application:	ELISA, SDS-PAGE (SDS), Western Blotting (WB), Crystallization (Crys)
Product Details	
Sequence:	NMLIALMGET VGQVSKESKH IWKLQWATTI LDIERSFPVF LRKAFRSGEM VTVGKSSDGT
	PDRRWCFRVD EVNWSHWNQN LGIINEDPGK NETYQYYGFS HTVGRLRRDR WSSVVPRVVE
	LNKNSNPDEV VVPLDSMGNP RCDGHQQGYP RKWRTDDAPL
	Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a
	special request, please contact us.
Characteristics:	<ul> <li>Made in Germany - from design to production - by highly experienced protein experts.</li> <li>Human TRPV4 Protein (raised in E. Coli) purified by multi-step, protein-specific process to ensure crystallization grade.</li> <li>State-of-the-art algorithm used for plasmid design (Gene synthesis).</li> </ul>
	This protein is a made to order protein and will be made for the first time for your order. Our
	experts in the lab will ensure that you receive a correctly folded protein.
	The big advantage of ordering our made-to-order proteins in comparison to ordering custom

made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).

When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered.

The concentration of our recombinant proteins is measured using the absorbance at 280nm.

The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.

The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

#### Purification:

Two step purification of proteins expressed in bacterial culture:

- 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.
- 2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

Purity:

>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.

Sterility:

0.22 µm filtered

Endotoxin Level:

Endotoxin has not been removed. Please contact us if you require endotoxin removal.

Grade:

Crystallography grade

#### **Target Details**

Target:	TRPV4
Alternative Name:	TRPV4 (TRPV4 Products)
Background:	Non-selective calcium permeant cation channel involved in osmotic sensitivity and
	mechanosensitivity. Activation by exposure to hypotonicity within the physiological range
	exhibits an outward rectification (PubMed:18826956, PubMed:18695040). Also activated by
	heat, low pH, citrate and phorbol esters (PubMed:18826956, PubMed:18695040). Increase of

intracellular Ca(2+) potentiates currents. Channel activity seems to be regulated by a calmodulin-dependent mechanism with a negative feedback mechanism (PubMed:12724311, PubMed:18826956). Promotes cell-cell junction formation in skin keratinocytes and plays an important role in the formation and/or maintenance of functional intercellular barriers (By similarity). Acts as a regulator of intracellular Ca(2+) in synoviocytes (PubMed:19759329). Plays an obligatory role as a molecular component in the nonselective cation channel activation induced by 4-alpha-phorbol 12,13-didecanoate and hypotonic stimulation in synoviocytes and also regulates production of IL-8 (PubMed:19759329). Together with PKD2, forms mechanoand thermosensitive channels in cilium (PubMed:18695040). Negatively regulates expression of PPARGC1A, UCP1, oxidative metabolism and respiration in adipocytes (By similarity). Regulates expression of chemokines and cytokines related to proinflammatory pathway in adipocytes (By similarity). Together with AQP5, controls regulatory volume decrease in salivary epithelial cells (By similarity). (ECO:0000250|UniProtKB:Q9EPK8, ECO:0000269|PubMed:11025659, ECO:0000269|PubMed:12724311, ECO:0000269|PubMed:18695040, ECO:0000269|PubMed:18826956, ECO:0000269|PubMed:19759329}.

Molecular Weight:	19.5 kDa Including tag.
UniProt:	Q9HBA0
Pathways: Hormone Transport, Cell-Cell Junction Organization	

### **Application Details**

Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a gurantee though.
Comment:	In cases in which it is highly likely that the recombinant protein with the default tag will be insoluble our protein lab may suggest a higher molecular weight tag (e.g. GST-tag) instead to increase solubility. We will discuss all possible options with you in detail to assure that you receive your protein of interest.
Restrictions:	For Research Use only

#### Handling

Format:	Liquid
Buffer:	100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer.

## Handling

Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	Unlimited (if stored properly)