antibodies .- online.com





USP44 Protein (AA 1-712) (His tag)





Go to Product page

Overview

Quantity:	1 mg
Target:	USP44
Protein Characteristics:	AA 1-712
Origin:	Human
Source:	Insect Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This USP44 protein is labelled with His tag.
Application:	ELISA, Western Blotting (WB), Crystallization (Crys), SDS-PAGE (SDS)

Product Details

Sequence:

MLAMDTCKHV GQLQLAQDHS SLNPQKWHCV DCNTTESIWA CLSCSHVACG RYIEEHALKH FQESSHPVAL EVNEMYVFCY LCDDYVLNDN TTGDLKLLRR TLSAIKSQNY HCTTRSGRFL RSMGTGDDSY FLHDGAQSLL QSEDQLYTAL WHRRRILMGK IFRTWFEQSP IGRKKQEEPF QEKIVVKREV KKRRQELEYQ VKAELESMPP RKSLRLQGLA QSTIIEIVSV QVPAQTPASP AKDKVLSTSE NEISQKVSDS SVKRRPIVTP GVTGLRNLGN TCYMNSVLQV LSHLLIFRQC FLKLDLNQWL AMTASEKTRS CKHPPVTDTV VYQMNECQEK DTGFVCSRQS SLSSGLSGGA SKGRKMELIQ PKEPTSQYIS LCHELHTLFQ VMWSGKWALV SPFAMLHSVW RLIPAFRGYA QQDAQEFLCE LLDKIQRELE TTGTSLPALI PTSQRKLIKQ VLNVVNNIFH GQLLSQVTCL ACDNKSNTIE PFWDLSLEFP ERYQCSGKDI ASQPCLVTEM LAKFTETEAL EGKIYVCDQC NSKRRRFSSK PVVLTEAQKQ LMICHLPQVL RLHLKRFRWS GRNNREKIGV HVGFEEILNM EPYCCRETLK SLRPECFIYD LSAVVMHHGK GFGSGHYTAY CYNSEGGFWV HCNDSKLSMC TMDEVCKAQA YILFYTQRVT ENGHSKLLPP ELLLGSQHPN EDADTSSNEI LS

Grade:

Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a special request, please contact us. Characteristics: Made in Germany - from design to production - by highly experienced protein experts. Human USP44 Protein (raised in Insect Cells) purified by multi-step, protein-specific process to ensure crystallization grade. • State-of-the-art algorithm used for plasmid design (Gene synthesis). This protein is a made to order protein and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein. The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified. In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization). When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered. The concentration of our recombinant proteins is measured using the absorbance at 280nm. The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer. The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein. Purification: Two step purification of proteins expressed in baculovirus infected SF9 insect cells: 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE. 2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot. >95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot. Purity: Sterility: 0.22 µm filtered Endotoxin Level: Protein is endotoxin free.

Crystallography grade

Target Details

Target:	USP44
Alternative Name:	USP44 (USP44 Products)
Background:	Deubiquitinase that plays a key regulatory role in the spindle assembly checkpoint or mitotic
	checkpoint by preventing premature anaphase onset. Acts by specifically mediating
	deubiquitination of CDC20, a negative regulator of the anaphase promoting
	complex/cyclosome (APC/C). Deubiquitination of CDC20 leads to stabilize the MAD2L1-CDC20
	APC/C ternary complex (also named mitotic checkpoint complex), thereby preventing
	premature activation of the APC/C. Promotes association of MAD2L1 with CDC20 and
	reinforces the spindle assembly checkpoint. Acts as a negative regulator of histone H2B
	(H2BK120ub1) ubiquitination. {ECO:0000269 PubMed:17443180,
	ECO:0000269 PubMed:22681888}.
Molecular Weight:	82.1 kDa Including tag.
UniProt:	Q9H0E7
Pathways:	M Phase
Application Details	
Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies
	as well. As the protein has not been tested for functional studies yet we cannot offer a gurantee
	though.
Comment:	In cases in which it is highly likely that the recombinant protein with the default tag will be
	insoluble our protein lab may suggest a higher molecular weight tag (e.g. GST-tag) instead to
	increase solubility. We will discuss all possible options with you in detail to assure that you
	receive your protein of interest.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.

Expiry Date:

Unlimited (if stored properly)

Images




Image 1. "Crystallography Grade" protein due to multi-step, protein-specific purification process