

# Datasheet for ABIN3096407 WEE2 Protein (AA 1-567) (Strep Tag)



Go to Product page

()	ve	r\/i	۱۸/
$\cup$	V C	1 / 1	 v v

Quantity:	250 μg
Target:	WEE2
Protein Characteristics:	AA 1-567
Origin:	Human
Source:	Cell-free protein synthesis (CFPS)
Protein Type:	Recombinant
Purification tag / Conjugate:	This WEE2 protein is labelled with Strep Tag.
Application:	ELISA, SDS-PAGE (SDS), Western Blotting (WB)

Product Details		
Brand:	AliCE®	
Sequence:	MDDKDIDKEL RQKLNFSYCE ETEIEGQKKV EESREASSQT PEKGEVQDSE AKGTPPWTPL	
	SNVHELDTSS EKDKESPDQI LRTPVSHPLK CPETPAQPDS RSKLLPSDSP STPKTMLSRL	
	VISPTGKLPS RGPKHLKLTP APLKDEMTSL ALVNINPFTP ESYKKLFLQS GGKRKIRGDL	
	EEAGPEEGKG GLPAKRCVLR ETNMASRYEK EFLEVEKIGV GEFGTVYKCI KRLDGCVYAI	
	KRSMKTFTEL SNENSALHEV YAHAVLGHHP HVVRYYSSWA EDDHMIIQNE YCNGGSLQAA	
	ISENTKSGNH FEEPKLKDIL LQISLGLNYI HNSSMVHLDI KPSNIFICHK MQSESSGVIE	
	EVENEADWFL SANVMYKIGD LGHATSINKP KVEEGDSRFL ANEILQEDYR HLPKADIFAL	
	GLTIAVAAGA ESLPTNGAAW HHIRKGNFPD VPQELSESFS SLLKNMIQPD AEQRPSAAAL	
	ARNTVLRPSL GKTEELQQQL NLEKFKTATL ERELREAQQA QSPQGYTHHG DTGVSGTHTG	
	SRSTKRLVGG KSARSSSFTS GEREPLH	
	Sequence without tag. The proposed Strep-Tag is based on experience s with the expressio	

## system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.

#### Characteristics:

#### Key Benefits:

- Made in Germany from design to production by highly experienced protein experts.
- · Protein expressed with ALiCE® and purified in one-step affinity chromatography
- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- · State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

#### Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require posttranslational modifications.
- During lysate production, the cell wall and other cellular components that are not required for
  protein production are removed, leaving only the protein production machinery and the
  mitochondria to drive the reaction. During our lysate completion steps, the additional
  components needed for protein production (amino acids, cofactors, etc.) are added to
  produce something that functions like a cell, but without the constraints of a living system all that's needed is the DNA that codes for the desired protein!

#### Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- · The protein's absorbance will be measured against its specific reference buffer.
- We use the Expasy's ProtParam tool to determine the absorption coefficient of each protein.

Purification:	One-step Strep-tag purification of proteins expressed in Almost Living Cell-Free Expression System (AliCE®).
Purity:	> 70-80 % as determined by SDS PAGE, Western Blot and analytical SEC (HPLC).
Grade:	custom-made

## **Target Details**

WEE2	
WEE2 (WEE2 Products)	
Wee1-like protein kinase 2 (EC 2.7.10.2) (Wee1-like protein kinase 1B) (Wee1B	
kinase),FUNCTION: Oocyte-specific protein tyrosine kinase that phosphorylates and inhibits	
CDK1/CDC2 and acts as a key regulator of meiosis during both prophase I and metaphase II	
(PubMed:29606300). Required to maintain meiotic arrest in oocytes during the germinal vesicle	
(GV) stage, a long period of quiescence at dictyate prophase I, by phosphorylating CDK1 at 'Tyr-	
15', leading to inhibit CDK1 activity and prevent meiotic reentry. Also required for metaphase II	
exit during egg activation by phosphorylating CDK1 at 'Tyr-15', to ensure exit from meiosis in	
oocytes and promote pronuclear formation (By similarity). {ECO:0000250 UniProtKB:Q66JT0,	
ECO:0000269 PubMed:29606300}.	
62.9 kDa	
P0C1S8	
M Phase	
In addition to the applications listed above we expect the protein to work for functional studies	
as well. As the protein has not been tested for functional studies yet we cannot offer a	
guarantee though.	
ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from	
Nicotiana tabacum c.v This contains all the protein expression machinery needed to produce	
even the most difficult-to-express proteins, including those that require post-translational	
modifications.	
During lysate production, the cell wall and other cellular components that are not required for	
protein production are removed, leaving only the protein production machinery and the	
mitochondria to drive the reaction. During our lysate completion steps, the additional	
mitochondria to drive the reaction. During our tysate completion steps, the additional	
components needed for protein production (amino acids, cofactors, etc.) are added to produce	
components needed for protein production (amino acids, cofactors, etc.) are added to produce	

## Handling

Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer.  Standard Storage Buffer: PBS pH 7.4, 10 % Glycerol <b>Might differ depending on protein.</b>
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	12 months