antibodies .- online.com





RNF144A Protein (AA 1-292) (Strep Tag)



Image



Go to Product page

Overview

Quantity:	1 mg
Target:	RNF144A
Protein Characteristics:	AA 1-292
Origin:	Human
Source:	Tobacco (Nicotiana tabacum)
Protein Type:	Recombinant
Purification tag / Conjugate:	This RNF144A protein is labelled with Strep Tag.
Application:	ELISA, Western Blotting (WB), SDS-PAGE (SDS)

Product Details

Sequence:

MTTTRYRPTW DLALDPLVSC KLCLGEYPVE QMTTIAQCQC IFCTLCLKQY VELLIKEGLE
TAISCPDAAC PKQGHLQENE IECMVAAEIM QRYKKLQFER EVLFDPCRTW CPASTCQAVC
QLQDVGLQTP QPVQCKACRM EFCSTCKASW HPGQGCPETM PITFLPGETS AAFKMEEDDA
PIKRCPKCKV YIERDEGCAQ MMCKNCKHAF CWYCLESLDD DFLLIHYDKG PCRNKLGHSR
ASVIWHRTQV VGIFAGFGLL LLVASPFLLL ATPFVLCCKC KCSKGDDDPL PT

Sequence without tag. The proposed Strep-Tag is based on experience s with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.

Characteristics:

Key Benefits:

- Made in Germany from design to production by highly experienced protein experts.
- Protein expressed with ALiCE® and purified by multi-step, protein-specific process to ensure correct folding and modification.

- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require posttranslational modifications.
- During lysate production, the cell wall and other cellular components that are not required for
 protein production are removed, leaving only the protein production machinery and the
 mitochondria to drive the reaction. During our lysate completion steps, the additional
 components needed for protein production (amino acids, cofactors, etc.) are added to
 produce something that functions like a cell, but without the constraints of a living system all that's needed is the DNA that codes for the desired protein!

Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.
- We use the Expasy's ProtParam tool to determine the absorption coefficient of each protein.

Purification:

Two step purification of proteins expressed in Almost Living Cell-Free Expression System (ALiCE®):

- 1. In a first purification step, the protein is purified from the cleared cell lysate using StrepTag capture material. Eluate fractions are analyzed by SDS-PAGE.
- 2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

Purity:

>80 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.

Endotoxin Level:

Low Endotoxin less than 1 EU/mg (< 0.1 ng/mg)

Product Details Grade: Crystallography grade Target Details Target: RNF144A Alternative Name RNF144A (RNF144A Products) Background: E3 ubiquitin-protein ligase RNF144A (EC 2.3.2.31) (RING finger protein 144A) (UbcM4interacting protein 4) (Ubiquitin-conjugating enzyme 7-interacting protein 4), FUNCTION: E3 ubiquitin-protein ligase which accepts ubiquitin from E2 ubiquitin-conjugating enzymes UBE2L3 and UBE2L6 in the form of a thioester and then directly transfers the ubiquitin to targeted substrates (PubMed:26216882). Mediates the ubiquitination and degradation of the DNA damage kinase PRKDC during DNA damage (PubMed:24979766). Positively regulates DNA virus or exogenous cytosolic DNA-triggered innate immune response by mediating STING1 ubiquitination and increasing its 'Lys-6'-linked ubiquitination and translocation from the endoplasmic reticulum to the Golgi leading to downstream signaling pathways (PubMed:37955227). Plays a positive role in EGF-dependent cell proliferation by prolonging EGF/EGFR signaling during EGF stimulation through EGFR ubiquitination (PubMed:30171075). Increases ERK activity independently of EGFR signaling by promoting polyubiquitination and subsequent degradation of VRK3 in the cytosol (PubMed:33067254). (ECO:0000250, ECO:0000269|PubMed:24979766, ECO:0000269|PubMed:26216882, ECO:0000269|PubMed:30171075, ECO:0000269|PubMed:33067254, ECO:0000269|PubMed:37955227}. Molecular Weight: 32.9 kDa UniProt: P50876 Application Details In addition to the applications listed above we expect the protein to work for functional studies **Application Notes:** as well. As the protein has not been tested for functional studies yet we cannot offer a

ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from

even the most difficult-to-express proteins, including those that require post-translational

Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce

During lysate production, the cell wall and other cellular components that are not required for

guarantee though.

modifications.

Comment:

protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

Restrictions:

For Research Use only

Handling

Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer. If you have a special request, please contact us.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	Unlimited (if stored properly)

Images



Image 1. "Crystallography Grade" protein due to multi-step, protein-specific purification process