# antibodies .- online.com





# HSD17B7 Protein (AA 1-341) (Strep Tag)



# **Image**



#### Overview

Quantity:	1 mg
Target:	HSD17B7
Protein Characteristics:	AA 1-341
Origin:	Human
Source:	Tobacco (Nicotiana tabacum)
Protein Type:	Recombinant
Purification tag / Conjugate:	This HSD17B7 protein is labelled with Strep Tag.
Application:	ELISA, Western Blotting (WB), SDS-PAGE (SDS)

#### **Product Details**

#### Sequence:

MRKVVLITGA SSGIGLALCK RLLAEDDELH LCLACRNMSK AEAVCAALLA SHPTAEVTIV QVDVSNLQSV FRASKELKQR FQRLDCIYLN AGIMPNPQLN IKALFFGLFS RKVIHMFSTA EGLLTQGDKI TADGLQEVFE TNVFGHFILI RELEPLLCHS DNPSQLIWTS SRSARKSNFS LEDFQHSKGK EPYSSSKYAT DLLSVALNRN FNQQGLYSNV ACPGTALTNL TYGILPPFIW TLLMPAILLL RFFANAFTLT PYNGTEALVW LFHQKPESLN PLIKYLSATT GFGRNYIMTQ KMDLDEDTAE KFYQKLLELE KHIRVTIQKT DNQARLSGSC L

Sequence without tag. The proposed Strep-Tag is based on experience s with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.

#### Characteristics:

Key Benefits:

- Made in Germany from design to production by highly experienced protein experts.
- · Protein expressed with ALiCE® and purified by multi-step, protein-specific process to ensure

correct folding and modification.

- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

#### Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require posttranslational modifications.
- During lysate production, the cell wall and other cellular components that are not required for
  protein production are removed, leaving only the protein production machinery and the
  mitochondria to drive the reaction. During our lysate completion steps, the additional
  components needed for protein production (amino acids, cofactors, etc.) are added to
  produce something that functions like a cell, but without the constraints of a living system all that's needed is the DNA that codes for the desired protein!

#### Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.
- We use the Expasy's ProtParam tool to determine the absorption coefficient of each protein.

#### Purification:

Two step purification of proteins expressed in Almost Living Cell-Free Expression System (ALiCE®):

- 1. In a first purification step, the protein is purified from the cleared cell lysate using StrepTag capture material. Eluate fractions are analyzed by SDS-PAGE.
- Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

Purity:

>80 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.

# **Product Details** Low Endotoxin less than 1 EU/mg (< 0.1 ng/mg) Endotoxin Level: Grade: Crystallography grade Target Details Target: HSD17B7 HSD17B7 (HSD17B7 Products) Alternative Name: Background: 3-keto-steroid reductase/17-beta-hydroxysteroid dehydrogenase 7 (17-beta-hydroxysteroid dehydrogenase 7) (17-beta-HSD 7) (3-keto-steroid reductase) (EC 1.1.1.270) (Dihydrotestosterone oxidoreductase) (EC 1.1.1.210) (Estradiol 17-beta-dehydrogenase 7) (EC 1.1.1.62) (Short chain dehydrogenase/reductase family 37C member 1), FUNCTION: Bifunctional enzyme involved in steroid-hormone metabolism and cholesterol biosynthesis (PubMed:12574203, PubMed:12732193, PubMed:12829805, PubMed:20659585, PubMed:19772289, PubMed:11165030). Catalyzes the NADP(H)-dependent reduction of estrogens and androgens and regulates the biological potency of these steroids. Converts estrone (E1) to a more potent estrogen, 17beta-estradiol (E2) (PubMed:12574203, PubMed:12732193, PubMed:19772289). Converts dihydrotestosterone (DHT) to its inactive form 5a-androstane-3b,17b-diol (PubMed:12574203, PubMed:12732193, PubMed:19772289). Converts moderately progesterone to 3beta-hydroxypregn-4-ene-20-one, leading to its inactivation (PubMed:12574203, PubMed:12732193). Additionally, participates in the postsqualene cholesterol biosynthesis, as a 3-ketosteroid reductase (PubMed:12829805, PubMed:20659585, PubMed:11165030). {ECO:0000269|PubMed:11165030, ECO:0000269|PubMed:12574203, ECO:0000269|PubMed:12732193, ECO:0000269|PubMed:12829805, ECO:0000269|PubMed:19772289, ECO:0000269|PubMed:20659585}., FUNCTION: [Isoform 3]: Does not have enzymatic activities toward E1 and DHT. {ECO:0000269|PubMed:12732193}. Molecular Weight: 38.2 kDa UniProt: P56937 Steroid Hormone Biosynthesis Pathways:

# **Application Details**

Application Notes:

In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.

# **Application Details**

#### Comment:

ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.

During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

Restrictions:

For Research Use only

# Handling

Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer. If you have a special request, please contact us.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	Unlimited (if stored properly)

#### **Images**



**Image 1.** "Crystallography Grade" protein due to multi-step, protein-specific purification process