
antibodies .- online.com





ST3GAL2 Protein (AA 1-350) (Strep Tag)





Go to Product page

Overview

Quantity:	1 mg
Target:	ST3GAL2
Protein Characteristics:	AA 1-350
Origin:	Human
Source:	Tobacco (Nicotiana tabacum)
Protein Type:	Recombinant
Purification tag / Conjugate:	This ST3GAL2 protein is labelled with Strep Tag.
Application:	ELISA, Western Blotting (WB), SDS-PAGE (SDS)

Product Details

Sequence:

MKCSLRVWFL SVAFLLVFIM SLLFTYSHHS MATLPYLDSG ALDGTHRVKL VPGYAGLQRL
SKERLSGKSC ACRRCMGDAG ASDWFDSHFD GNISPVWTRE NMDLPPDVQR WWMMLQPQFK
SHNTNEVLEK LFQIVPGENP YRFRDPHQCR RCAVVGNSGN LRGSGYGQDV DGHNFIMRMN
QAPTVGFEQD VGSRTTHHFM YPESAKNLPA NVSFVLVPFK VLDLLWIASA LSTGQIRFTY
APVKSFLRVD KEKVQIYNPA FFKYIHDRWT EHHGRYPSTG MLVLFFALHV CDEVNVYGFG
ADSRGNWHHY WENNRYAGEF RKTGVHDADF EAHIIDMLAK ASKIEVYRGN

Sequence without tag. The proposed Strep-Tag is based on experience s with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.

Characteristics:

Key Benefits:

- Made in Germany from design to production by highly experienced protein experts.
- · Protein expressed with ALiCE® and purified by multi-step, protein-specific process to ensure

correct folding and modification.

- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require posttranslational modifications.
- During lysate production, the cell wall and other cellular components that are not required for
 protein production are removed, leaving only the protein production machinery and the
 mitochondria to drive the reaction. During our lysate completion steps, the additional
 components needed for protein production (amino acids, cofactors, etc.) are added to
 produce something that functions like a cell, but without the constraints of a living system all that's needed is the DNA that codes for the desired protein!

Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.
- We use the Expasy's ProtParam tool to determine the absorption coefficient of each protein.

Purification:

Two step purification of proteins expressed in Almost Living Cell-Free Expression System (ALiCE®):

- 1. In a first purification step, the protein is purified from the cleared cell lysate using StrepTag capture material. Eluate fractions are analyzed by SDS-PAGE.
- Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

Purity:

>80 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.

Product Details Low Endotoxin less than 1 EU/mg (< 0.1 ng/mg) Endotoxin Level: Grade: Crystallography grade Target Details Target: ST3GAL2 Alternative Name: ST3GAL2 (ST3GAL2 Products) Background: CMP-N-acetylneuraminate-beta-galactosamide-alpha-2,3-sialyltransferase 2 (Alpha 2,3-ST 2) (Beta-galactoside alpha-2,3-sialyltransferase 2) (EC 2.4.3.4) (Gal-NAc6S) (Gal-beta-1,3-GalNAcalpha-2,3-sialyltransferase) (Monosialoganglioside sialyltransferase) (EC 2.4.3.2) (ST3Gal II) (ST3GalII) (ST3GalA.2) (Sialyltransferase 4B) (SIAT4-B),FUNCTION: A beta-galactoside alpha2-3 sialyltransferase primarily involved in terminal sialylation of ganglio and globo series glycolipids (PubMed:8920913, PubMed:9266697). Catalyzes the transfer of sialic acid (N-acetyl-neuraminic acid, Neu5Ac) from the nucleotide sugar donor CMP-Neu5Ac onto acceptor Galbeta-(1->3)-GalNAc-terminated glycoconjugates through an alpha2-3 linkage (PubMed:8920913, PubMed:9266697, PubMed:25916169). Sialylates GM1/GM1a, GA1/asialo-GM1 and GD1b gangliosides to form GD1a, GM1b and GT1b, respectively (PubMed:8920913, PubMed:9266697). Together with ST3GAL3, primarily responsible for biosynthesis of brain GD1a and GT1b that function as ligands for myelin-associated glycoprotein MAG on axons, regulating MAG expression and axonal myelin stability and regeneration (By similarity). Via GT1b regulates TLR2 signaling in spinal cord microglia in response to nerve injury (By similarity). Responsible for the sialylation of the pluripotent stem cell- and cancer stem cellassociated antigen SSEA3, forming SSEA4 (PubMed:12716912). Sialylates with low efficiency asialofetuin, presumably onto O-glycosidically linked Galbeta-(1->3)-GalNAc-O-Ser (PubMed:9266697, PubMed:25916169). {ECO:0000250|UniProtKB:Q11204, ECO:0000269|PubMed:12716912, ECO:0000269|PubMed:25916169, ECO:0000269|PubMed:8920913, ECO:0000269|PubMed:9266697}. Molecular Weight: 40.2 kDa UniProt: Q16842 Pathways: Glycosaminoglycan Metabolic Process **Application Details**

Application Notes:

In addition to the applications listed above we expect the protein to work for functional studies

as well. As the protein has not been tested for functional studies yet we cannot offer a

Application Details

, ipplication betains	
	guarantee though.
Comment:	ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from
	Nicotiana tabacum c.v This contains all the protein expression machinery needed to produce
	even the most difficult-to-express proteins, including those that require post-translational
	modifications.
	During lysate production, the cell wall and other cellular components that are not required for
	protein production are removed, leaving only the protein production machinery and the
	mitochondria to drive the reaction. During our lysate completion steps, the additional
	components needed for protein production (amino acids, cofactors, etc.) are added to produce
	something that functions like a cell, but without the constraints of a living system - all that's
	needed is the DNA that codes for the desired protein!
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer. If you have a special request,
	please contact us.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	Unlimited (if stored properly)




Image 1. "Crystallography Grade" protein due to multi-step, protein-specific purification process