# antibodies .- online.com





# SLC25A29 Protein (AA 1-303) (Strep Tag)





Go to Product page

#### Overview

Quantity:	1 mg
Target:	SLC25A29
Protein Characteristics:	AA 1-303
Origin:	Human
Source:	Tobacco (Nicotiana tabacum)
Protein Type:	Recombinant
Purification tag / Conjugate:	This SLC25A29 protein is labelled with Strep Tag.
Application:	ELISA, SDS-PAGE (SDS), Western Blotting (WB)

#### **Product Details**

#### Sequence:

MALDFLAGCA GGVAGVLVGH PFDTVKVRLQ VQSVEKPQYR GTLHCFKSII KQESVLGLYK
GLGSPLMGLT FINALVFGVQ GNTLRALGHD SPLNQFLAGA AAGAIQCVIC CPMELAKTRL
QLQDAGPART YKGSLDCLAQ IYGHEGLRGV NRGMVSTLLR ETPSFGVYFL TYDALTRALG
CEPGDRLLVP KLLLAGGTSG IVSWLSTYPV DVVKSRLQAD GLRGAPRYRG ILDCVHQSYR
AEGWRVFTRG LASTLLRAFP VNAATFATVT VVLTYARGEE AGPEGEAVPA APAGPALAQP SSL

Sequence without tag. The proposed Strep-Tag is based on experience s with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.

#### Characteristics:

Key Benefits:

- Made in Germany from design to production by highly experienced protein experts.
- Protein expressed with ALiCE® and purified by multi-step, protein-specific process to ensure correct folding and modification.

- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

#### Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require posttranslational modifications.
- During lysate production, the cell wall and other cellular components that are not required for
  protein production are removed, leaving only the protein production machinery and the
  mitochondria to drive the reaction. During our lysate completion steps, the additional
  components needed for protein production (amino acids, cofactors, etc.) are added to
  produce something that functions like a cell, but without the constraints of a living system all that's needed is the DNA that codes for the desired protein!

#### Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.
- We use the Expasy's ProtParam tool to determine the absorption coefficient of each protein.

#### Purification:

Two step purification of proteins expressed in Almost Living Cell-Free Expression System (ALiCE®):

- 1. In a first purification step, the protein is purified from the cleared cell lysate using StrepTag capture material. Eluate fractions are analyzed by SDS-PAGE.
- Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

Purity:

>80 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.

Endotoxin Level:

Low Endotoxin less than 1 EU/mg (< 0.1 ng/mg)

# **Product Details** Grade: Crystallography grade **Target Details** Target: SLC25A29 Alternative Name SLC25A29 (SLC25A29 Products) Background: Mitochondrial basic amino acids transporter (Carnitine/acylcarnitine translocase-like) (CACTlike) (Mitochondrial carnitine/acylcarnitine carrier protein CACL) (Mitochondrial ornithine transporter 3) (Solute carrier family 25 member 29), FUNCTION: Mitochondrial transporter of arginine, lysine, homoarginine, methylarginine and, to a much lesser extent, ornithine and histidine (PubMed:19287344, PubMed:24652292). Does not transport carnitine nor acylcarnitines (PubMed:24652292). Functions by both counter-exchange and uniport mechanisms (PubMed:24652292). Plays a physiological role in the import of basic amino acids into mitochondria for mitochondrial protein synthesis and amino acid degradation (PubMed:19287344, PubMed:24652292). {ECO:0000269|PubMed:19287344, ECO:0000269|PubMed:24652292}. Molecular Weight: 32.1 kDa UniProt: **Q8N8R3** Application Details **Application Notes:** In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though. Comment: ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications. During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the

Restrictions: For Research Use only

needed is the DNA that codes for the desired protein!

mitochondria to drive the reaction. During our lysate completion steps, the additional


components needed for protein production (amino acids, cofactors, etc.) are added to produce

something that functions like a cell, but without the constraints of a living system - all that's

### Handling

Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer. If you have a special request, please contact us.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	Unlimited (if stored properly)

## Images



**Image 1.** "Crystallography Grade" protein due to multi-step, protein-specific purification process