antibodies .- online.com





C4orf49 Protein (AA 1-240) (Strep Tag)



Image



Go to Product pag

Overview

Quantity:	1 mg
Target:	C4orf49
Protein Characteristics:	AA 1-240
Origin:	Human
Source:	Tobacco (Nicotiana tabacum)
Protein Type:	Recombinant
Purification tag / Conjugate:	This C4orf49 protein is labelled with Strep Tag.
Application:	ELISA, Western Blotting (WB), SDS-PAGE (SDS)

Product Details

Sequ	ience:
------	--------

MYLRRAVSKT LALPLRAPPN PAPLGKDASL RRMSSNRFPG SSGSNMIYYL VVGVTVSAGG
YYAYKTVTSD QAKHTEHKTN LKEKTKAEIH PFQGEKENVA ETEKASSEAP EELIVEAEVV
DAEESPSATV VVIKEASACP GHVEAAPETT AVSAETGPEV TDAAARETTE VNPETTPEVT
NAALDEAVTI DNDKDTTKNE TSDEYAELEE ENSPAESESS AGDDLQEEAS VGSEAASAQG

Sequence without tag. The proposed Strep-Tag is based on experience s with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.

Characteristics:

Key Benefits:

- Made in Germany from design to production by highly experienced protein experts.
- Protein expressed with ALiCE® and purified by multi-step, protein-specific process to ensure correct folding and modification.
- · These proteins are normally active (enzymatically functional) as our customers have

reported (not tested by us and not guaranteed).

• State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require posttranslational modifications.
- During lysate production, the cell wall and other cellular components that are not required for
 protein production are removed, leaving only the protein production machinery and the
 mitochondria to drive the reaction. During our lysate completion steps, the additional
 components needed for protein production (amino acids, cofactors, etc.) are added to
 produce something that functions like a cell, but without the constraints of a living system all that's needed is the DNA that codes for the desired protein!

Concentration:

- · The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.
- · We use the Expasy's ProtParam tool to determine the absorption coefficient of each protein.

Purification:	Two step purification of proteins expressed in Almost Living Cell-Free Expression System
	(ALiCE®):
	1. In a first purification step, the protein is purified from the cleared cell lysate using StrepTag capture material. Eluate fractions are analyzed by SDS-PAGE.
	Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.
Purity:	>80 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.
Endotoxin Level:	Low Endotoxin less than 1 EU/mg (< 0.1 ng/mg)

Product Details Grade: Crystallography grade **Target Details** Target: C4orf49 Alternative Name MGARP (C4orf49 Products) Background: Protein MGARP (Corneal endothelium-specific protein 1) (CESP-1) (Hypoxia up-regulated mitochondrial movement regulator protein) (Mitochondria-localized glutamic acid-rich protein) (Ovary-specific acidic protein), FUNCTION: Plays a role in the trafficking of mitochondria along microtubules. Regulates the kinesin-mediated axonal transport of mitochondria to nerve terminals along microtubules during hypoxia. Participates in the translocation of TRAK2/GRIF1 from the cytoplasm to the mitochondrion. Also plays a role in steroidogenesis through maintenance of mitochondrial abundance and morphology (By similarity). Plays an inhibitory role during neocortex development by regulating mitochondrial morphology, distribution and motility in neocortical neurons (By similarity). {ECO:0000250|UniProtKB:Q8VI64}. Molecular Weight: 25.4 kDa UniProt: Q8TDB4 **Application Details Application Notes:** In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though. Comment: ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications. During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

For Research Use only

Restrictions:

Handling

Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer. If you have a special request, please contact us.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	Unlimited (if stored properly)

Images




Image 1. "Crystallography Grade" protein due to multi-step, protein-specific purification process