antibodies

## Datasheet for ABIN3108247 Endomucin Protein (EMCN) (AA 19-261) (rho-1D4 tag)



Image

Overview	
Quantity:	1 mg
Target:	Endomucin (EMCN)
Protein Characteristics:	AA 19-261
Origin:	Human
Source:	Insect Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This Endomucin protein is labelled with rho-1D4 tag.
Application:	SDS-PAGE (SDS), ELISA, Western Blotting (WB), Crystallization (Crys)

## Product Details

Sequence:	NSTGVLEAAN NSLVVTTTKP SITTPNTESL QKNVVTPTTG TTPKGTITNE LLKMSLMSTA
	TFLTSKDEGL KATTTDVRKN DSIISNVTVT SVTLPNAVST LQSSKPKTET QSSIKTTEIP
	GSVLQPDASP SKTGTLTSIP VTIPENTSQS QVIGTEGGKN ASTSATSRSY SSIILPVVIA LIVITLSVFV
	LVGLYRMCWK ADPGTPENGN DQPQSDKESV KLLTVKTISH ESGEHSAQGK TKN
	Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a
	special request, please contact us.
Characteristics:	<ul> <li>Made in Germany - from design to production - by highly experienced protein experts.</li> <li>Human EMCN Protein (raised in Insect Cells) purified by multi-step, protein-specific process</li> </ul>
	<ul><li>to ensure crystallization grade.</li><li>State-of-the-art algorithm used for plasmid design (Gene synthesis).</li></ul>
	State-or-the-art algorithm used for plasmid design (Gene Synthesis).
	This protein is a made to order protein and will be made for the first time for your order. Our
	experts in the lab will ensure that you receive a correctly folded protein.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/4 | Product datasheet for ABIN3108247 | 09/11/2023 | Copyright antibodies-online. All rights reserved.

made proteins from other companies is to cannot be expressed or purified.In the unlikely event that the protein can (other companies might charge you for a custom-made proteins, e.g. fees might a experiments or purification optimization)When you order this made-to-order protect folded protein. With no financial risk on y protein experts will do everything to mak The concentration of our recombinant pr The protein's absorbance will be measure specific reference buffer.Purification:Three step purification of membrane pro cells: 1. Membrane proteins are fractioned by the	to-order proteins in comparison to ordering custom hat there is no financial obligation in case the protein of be expressed or purified we do not charge anything ny performed steps in the expression process for oply for the expression plasmid, the first expression on you will only pay upon receival of the correctly our end you can rest assured that our experienced e sure that you receive the protein you ordered. ofteins is measured using the absorbance at 280nm. ed in several dilutions and is measured against its ated using its specific absorption coefficient. We use e the absorption coefficient of each protein. teins expressed in baculovirus infected SF9 insect
cannot be expressed or purified.In the unlikely event that the protein canner(other companies might charge you for a custom-made proteins, e.g. fees might a experiments or purification optimization)When you order this made-to-order protein folded protein. With no financial risk on y protein experts will do everything to make The concentration of our recombinant pri The protein's absorbance will be measure specific reference buffer.Purification:Three step purification of membrane pro cells: 1. Membrane proteins are fractioned by the	ot be expressed or purified we do not charge anything ny performed steps in the expression process for oply for the expression plasmid, the first expression in you will only pay upon receival of the correctly our end you can rest assured that our experienced e sure that you receive the protein you ordered. oteins is measured using the absorbance at 280nm. ed in several dilutions and is measured against its ated using its specific absorption coefficient. We use e the absorption coefficient of each protein.
In the unlikely event that the protein cannot (other companies might charge you for a custom-made proteins, e.g. fees might and experiments or purification optimization) When you order this made-to-order protein folded protein. With no financial risk on y protein experts will do everything to make The concentration of our recombinant protein's absorbance will be measured specific reference buffer. The concentration of the protein is calculated the Expasy's protparam tool to determined the Expasy's protparam tool to determined the Expasy's proteins are fractioned by the first or the proteins are fractioned by the experiments or the proteins are fractioned by the experiments are fractioned by the first or the proteins are fracted by the first or the proteins are fracted by the first or the proteins are fr	ny performed steps in the expression process for oply for the expression plasmid, the first expression in you will only pay upon receival of the correctly our end you can rest assured that our experienced e sure that you receive the protein you ordered. oteins is measured using the absorbance at 280nm. ed in several dilutions and is measured against its ated using its specific absorption coefficient. We use e the absorption coefficient of each protein.
(other companies might charge you for a custom-made proteins, e.g. fees might a experiments or purification optimization) When you order this made-to-order protect folded protein. With no financial risk on y protein experts will do everything to mak The concentration of our recombinant pr The protein's absorbance will be measure specific reference buffer.The concentration of the protein is calcu the Expasy's protparam tool to determinePurification:Three step purification of membrane pro cells: 1. Membrane proteins are fractioned by the	ny performed steps in the expression process for oply for the expression plasmid, the first expression in you will only pay upon receival of the correctly our end you can rest assured that our experienced e sure that you receive the protein you ordered. oteins is measured using the absorbance at 280nm. ed in several dilutions and is measured against its ated using its specific absorption coefficient. We use e the absorption coefficient of each protein.
custom-made proteins, e.g. fees might a experiments or purification optimization) When you order this made-to-order prote folded protein. With no financial risk on y protein experts will do everything to mak The concentration of our recombinant pr The protein's absorbance will be measure specific reference buffer. The concentration of the protein is calcu the Expasy's protparam tool to determine Purification:Purification:Three step purification of membrane pro cells: 1. Membrane proteins are fractioned by the	pply for the expression plasmid, the first expression in you will only pay upon receival of the correctly our end you can rest assured that our experienced e sure that you receive the protein you ordered. oteins is measured using the absorbance at 280nm. ed in several dilutions and is measured against its ated using its specific absorption coefficient. We use e the absorption coefficient of each protein.
experiments or purification optimization)When you order this made-to-order protectfolded protein. With no financial risk on yprotein experts will do everything to makeThe concentration of our recombinant prThe protein's absorbance will be measurespecific reference buffer.The concentration of the protein is calculthe Expasy's protparam tool to determinePurification:Three step purification of membrane process1. Membrane proteins are fractioned by the	in you will only pay upon receival of the correctly our end you can rest assured that our experienced e sure that you receive the protein you ordered. oteins is measured using the absorbance at 280nm. ed in several dilutions and is measured against its ated using its specific absorption coefficient. We use e the absorption coefficient of each protein.
When you order this made-to-order protectfolded protein. With no financial risk on yprotein experts will do everything to makThe concentration of our recombinant prThe protein's absorbance will be measuredspecific reference buffer.The concentration of the protein is calculatedthe Expasy's protparam tool to determinedPurification:Three step purification of membrane process1. Membrane proteins are fractioned by the step of the protein is calculated	in you will only pay upon receival of the correctly our end you can rest assured that our experienced e sure that you receive the protein you ordered. oteins is measured using the absorbance at 280nm. ed in several dilutions and is measured against its ated using its specific absorption coefficient. We use e the absorption coefficient of each protein.
folded protein. With no financial risk on y protein experts will do everything to mak The concentration of our recombinant pr The protein's absorbance will be measure specific reference buffer.The concentration of the protein is calcu the Expasy's protparam tool to determinePurification:Three step purification of membrane pro cells: 1. Membrane proteins are fractioned by the function of the protein is calculed by the function of the protein of the protein of the protein is calculed by the function of the protein	our end you can rest assured that our experienced e sure that you receive the protein you ordered. oteins is measured using the absorbance at 280nm. ed in several dilutions and is measured against its ated using its specific absorption coefficient. We use e the absorption coefficient of each protein.
protein experts will do everything to makThe concentration of our recombinant prThe protein's absorbance will be measurespecific reference buffer.The concentration of the protein is calcuthe Expasy's protparam tool to determinePurification:Three step purification of membrane procells:1. Membrane proteins are fractioned by the	e sure that you receive the protein you ordered. oteins is measured using the absorbance at 280nm. ed in several dilutions and is measured against its ated using its specific absorption coefficient. We use the absorption coefficient of each protein.
The concentration of our recombinant pr         The protein's absorbance will be measure         specific reference buffer.         The concentration of the protein is calcu         the Expasy's protparam tool to determine         Purification:       Three step purification of membrane pro         cells:         1. Membrane proteins are fractioned by the step of the step in the step of the ste	oteins is measured using the absorbance at 280nm. ed in several dilutions and is measured against its ated using its specific absorption coefficient. We use the absorption coefficient of each protein.
The protein's absorbance will be measure specific reference buffer.The concentration of the protein is calcu the Expasy's protparam tool to determinePurification:Three step purification of membrane pro cells: 1. Membrane proteins are fractioned by the	ed in several dilutions and is measured against its ated using its specific absorption coefficient. We use the absorption coefficient of each protein.
specific reference buffer.         The concentration of the protein is calcu         the Expasy's protparam tool to determine         Purification:       Three step purification of membrane procells:         1. Membrane proteins are fractioned by the step purification of the protein step fractioned by the step purification of the protein step fractioned by the step purification of the protein step fractioned by the step purification of the protein step fractioned by the step purification of the protein step fractioned by the step purification of the protein step fractioned by the step purification of the protein step fractioned by the step purification of the protein step fractioned by the step purification of the protein step fractioned by the step purification of the protein step fractioned by the step purification of the protein step fractioned by the step purification of the protein step fractioned by the step purification of the protein step fractioned by the step purification of the protein step fractioned by the step purification of the protein step fractioned by the step purification of the protein step fractioned by the step purification of the protein step purification of the prot	ated using its specific absorption coefficient. We use the absorption coefficient of each protein.
The concentration of the protein is calcu         the Expasy's protparam tool to determine         Purification:       Three step purification of membrane procedules         cells:       1. Membrane proteins are fractioned by the step purification of the protein step fractioned by the step purification of the protein step fractioned by the step purification of the protein step fractioned by the step purification of the protein step fractioned by the step purification of the protein step fractioned by the step purification of the protein step fractioned by the step purification of the protein step fractioned by the step purification of the protein step fractioned by the step purification of the protein step fractioned by the step purification of the protein step fractioned by the step purification of the protein step fractioned by the step purification of the protein step fractioned by the step purification of the protein step fractioned by the step purification of the protein step fractioned by the step purification of the protein step fractioned by the step purification of the protein step fractioned by the step purification of the protein s	e the absorption coefficient of each protein.
the Expasy's protparam tool to determine         Purification:       Three step purification of membrane procedure         cells:       1. Membrane proteins are fractioned by the step purification of the step purificatio	e the absorption coefficient of each protein.
Purification:       Three step purification of membrane process         cells:       1. Membrane proteins are fractioned by the step purification of the step purification	
cells: 1. Membrane proteins are fractioned by t	teins expressed in baculovirus infected SF9 insect
1. Membrane proteins are fractioned by u	
different detergents (detergent screen	Iltracentrifugation and subsequently solubilized with
	. Samples are analyzed by Western blot.
	for solubilization and the proteins are purified via their plumns: one DTT resistant, the other one not. Eluate
fractions are analyzed by Western blot	
-	t purification are subjected to second purification step . Eluate fractions are analyzed by SDS-PAGE and
Purity: >95 % as determined by SDS PAGE, Size	Exclusion Chromatography and Western Blot.
Sterility: 0.22 µm filtered	
Endotoxin Level: Protein is endotoxin-free.	
Grade: Crystallography grade	
Target Details	
Target: Endomucin (EMCN)	
Alternative Name: EMCN (EMCN Products)	
Background: Endothelial sialomucin, also called endor	nucin or mucin-like sialoglycoprotain which interferes

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/4 | Product datasheet for ABIN3108247 | 09/11/2023 | Copyright antibodies-online. All rights reserved.

Target Details	
	with the assembly of focal adhesion complexes and inhibits interaction between cells and the extracellular matrix.
Molecular Weight:	26.6 kDa Including tag.
UniProt:	Q9ULC0
Application Details	
Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.
Comment:	In cases in which it is highly likely that the recombinant protein with the default tag will be insoluble our protein lab may suggest a higher molecular weight tag (e.g. GST-tag) instead to increase solubility. We will discuss all possible options with you in detail to assure that you receive your protein of interest.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	Unlimited (if stored properly)



**Image 1.** "Crystallography Grade" protein due to multi-step, protein-specific purification process

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 4/4 | Product datasheet for ABIN3108247 | 09/11/2023 | Copyright antibodies-online. All rights reserved.