Datasheet for ABIN3110347

FADS1 Protein (AA 1-444) (Strep Tag)

Go to Product page

Overview

Quantity:	250 µg
Target:	FADS1
Protein Characteristics:	AA 1-444
Origin:	Human
Source:	Cell-free protein synthesis (CFPS)
Protein Type:	Recombinant
Purification tag / Conjugate:	This FADS1 protein is labelled with Strep Tag.
Application:	Western Blotting (WB), SDS-PAGE (SDS), ELISA

Product Details

Brand:	AliCE®
Sequence:	MAPDPVAAET AAQGPTPRYF TWDEVAQRSG CEERWLVIDR KVYNISEFTR RHPGGSRVIS HYAGQDATDP FVAFHINKGL VKKYMNSLLI GELSPEQPSF EPTKNKELTD EFRELRATVE RMGLMKANHV FFLLYLLHIL LLDGAAWLTL WVFGTSFLPF LLCAVLLSAV QAQAGWLQHD FGHLSVFSTS KWNHLLHHFV IGHLKGAPAS WWNHMHFQHH AKPNCFRKDP DINMHPFFFA LGKILSVELG KQKKKYMPYN HQHKYFFLIG PPALLPLYFQ WYIFYFVIQR KKWVDLAWMI TFYVRFFLTY VPLLGLKAFL GLFFIVRFLE SNWFVWVTQM NHIPMHIDHD RNMDWVSTQL QATCNVHKSA FNDWFSGHLN FQIEHHLFPT MPRHNYHKVA PLVQSLCAKH GIEYQSKPLL SAFADIIHSL KESGQLWLDA YLHQ **Sequence without tag. The proposed Strep-Tag is based on experience s with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.**

Product Details

Characteristics:	Key Benefits: • Made in Germany - from design to production - by highly experienced protein experts. • Protein expressed with ALiCE® and purified in one-step affinity chromatography • These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed). • State-of-the-art algorithm used for plasmid design (Gene synthesis). This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein. The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified. Expression System: • ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications. • During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein! Concentration: • The concentration of our recombinant proteins is measured using the absorbance at 280nm. • The protein's absorbance will be measured against its specific reference buffer. • We use the Expasy's ProtParam tool to determine the absorption coefficient of each protein.
Purification:	One-step Strep-tag purification of proteins expressed in Almost Living Cell-Free Expression System (AliCE®).
Purity:	> 70-80 % as determined by SDS PAGE, Western Blot and analytical SEC (HPLC).
Grade:	custom-made

Target Details

Target:	FADS1
Alternative Name:	FADS1 (FADS1 Products)
Background:	Acyl-CoA (8-3)-desaturase (EC 1.14.19.44) (Delta(5) fatty acid desaturase) (D5D) (Delta(5) desaturase) (Delta-5 desaturase) (Fatty acid desaturase 1),FUNCTION: [Isoform 1]: Acts as a front-end fatty acyl-coenzyme A (CoA) desaturase that introduces a cis double bond at carbon 5 located between a preexisting double bond and the carboxyl end of the fatty acyl chain. Involved in biosynthesis of highly unsaturated fatty acids (HUFA) from the essential polyunsaturated fatty acids (PUFA) linoleic acid (LA) (18:2n-6) and alpha-linolenic acid (ALA) (18:3n-3) precursors. Specifically, desaturates dihomo-gamma-linoleoate (DGLA) (20:3n-6) and eicosatetraenoate (ETA) (20:4n-3) to generate arachidonate (AA) (20:4n-6) and eicosapentaenoate (EPA) (20:5n-3), respectively (PubMed:10601301, PubMed:10769175). As a rate limiting enzyme for DGLA (20:3n-6) and AA (20:4n-6)-derived eicosanoid biosynthesis, controls the metabolism of inflammatory lipids like prostaglandin E2, critical for efficient acute inflammatory response and maintenance of epithelium homeostasis. Contributes to membrane phospholipid biosynthesis by providing AA (20:4n-6) as a major acyl chain esterified into phospholipids. In particular, regulates phosphatidylinositol-4,5-bisphosphate levels, modulating inflammatory cytokine production in T-cells (By similarity). Also desaturates (11E)-octadecenoate (trans-vaccenoate)(18:1n-9), a metabolite in the biohydrogenation pathway of LA (18:2n-6) (By similarity). {ECO:0000250|UniProtKB:Q920L1, ECO:0000250|UniProtKB:Q920R3, ECO:0000269|PubMed:10601301, ECO:0000269|PubMed:10769175}., FUNCTION: [Isoform 2]: Does not exhibit any catalytic activity toward 20:3n-6, but it may enhance FADS2 activity. {ECO:0000250|UniProtKB:A4UVI1}.
Molecular Weight:	52.0 kDa
UniProt:	O60427
Pathways:	Regulation of Lipid Metabolism by PPARalpha

Application Details

Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.
Comment:	ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational

Application Details

	modifications. During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!
Restrictions:	For Research Use only

Handling

Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer. Standard Storage Buffer: PBS pH 7.4, 10 % Glycerol **Might differ depending on protein.**
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	12 months